

Datasheet: MCA2344

BATCH NUMBER 174191

Description:	MOUSE ANTI HUMAN CD221
Specificity:	CD221
Other names:	IGF-1 RECEPTOR
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	1H7
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10
Immunohistology - Frozen			▪	
Immunohistology - Paraffin (1)	▪			1/10 - 1/40
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires protein digestion pre-treatment of paraffin sections e.g. trypsin or pronase.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline

Preservative Stabilisers	0.09% sodium azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Purified human placental IGF-I receptor.
External Database Links	<p>UniProt: P08069 Related reagents</p> <p>Entrez Gene: 3480 IGF1R Related reagents</p>
RRID	AB_566647
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the SP2/0 myeloma cell line.
Specificity	<p>Mouse anti Human CD221 antibody, clone 1H7 recognizes human CD221, a ~155 kDa receptor tyrosine kinase, also known as Insulin-like growth factor I receptor (IGF-I Receptor). CD221 is composed of two extracellular alpha-subunits and two transmembrane beta-subunits. Clone 1H7 recognizes an epitope in the alpha subunits of CD221, demonstrated by western blotting (Li et al.1993).</p> <p>CD221 is expressed in a range of tissues, including kidney, liver, placenta, mammary gland, brain, ovary and skin.</p> <p>The ligands for CD221 include IGF-I and IGF-II, which bind to CD221 and activate tyrosine kinase activity, resulting in phosphorylation of several intracellular signaling proteins. Clone 1H7 is reported to partially block binding of IGF-I and IGF-II to CD221 (Li et al.1993).</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
Histology Positive Control Tissue	Human pancreas or placenta
References	<ol style="list-style-type: none"> Li, S.L. <i>et al.</i> (1993) Two new monoclonal antibodies against the alpha subunit of the human insulin-like growth factor-I receptor. Biochem Biophys Res Commun. 196 (1): 92-8. Beauvais, D.M. & Rapraeger, A.C. (2010) Syndecan-1 couples the insulin-like growth factor-1 receptor to inside-out integrin activation. J Cell Sci. 123 (Pt 21): 3796-807. Krieger, C.C. <i>et al.</i> (2016) Thyrotropin/IGF-1 receptor crosstalk in Graves' ophthalmopathy pathogenesis. J Clin Endocrinol Metab. Apr 4: jc20161315. [Epub ahead of print] van Valen, F. <i>et al.</i> (2012) A Novel Role of IGF1 in Apo2L/TRAIL-Mediated Apoptosis of

Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2344
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Rabbit Anti Mouse IgG (STAR12...)	RPE
Rabbit Anti Mouse IgG (STAR13...)	HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M450874:260616'

Printed on 16 Jun 2026