

Datasheet: MCA2344PE BATCH NUMBER 167043

Description:	MOUSE ANTI HUMAN CD221:RPE
Specificity:	CD221
Other names:	IGF-1 RECEPTOR
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	1H7
lsotype:	lgG1
Quantity:	100 TESTS

Product Details

Applications	derived from testing wi communications from	ithin our laboratories, the originators. Pleas al protocol recomme	he following application peer-reviewed publica refer to references in ndations, please visit <u>w</u>	tions or personal dicated for further		
		Yes No	Not Determined	Suggested Dilution		
	Flow Cytometry	•		Neat		
	necessarily exclude its	s use in such procedu mmended that the us	er titrates the product f	g dilutions are given as		
Target Species	Human					
Product Form	Purified IgG conjugate	d to R. Phycoerythrir	n (RPE) - lyophilized			
Reconstitution	Reconstitute with 1 ml	distilled water				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)			
	RPE 488nm laser	496	578			
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant					
Buffer Solution	Phosphate buffered sa	lline				
Preservative Stabilisers	0.09% sodium azide (1 1% bovine serum albu	0)				

	5% sucrose
Immunogen	Purified human placental IGF-I receptor.
External Database Links	UniProt: <u>P08069</u> <u>Related reagents</u> Entrez Gene: <u>3480</u> IGF1R <u>Related reagents</u>
RRID	AB_566646
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the SP2/0 myeloma cell line.
Specificity	 Mouse anti Human CD221 antibody, clone 1H7 recognizes human CD221, a ~155 kDa receptor tyrosine kinase, also known as Insulin-like growth factor I receptor (IGF-I Receptor). CD221 is composed of two extracellular alpha-subunits and two transmembrane beta-subunits. Clone 1H7 recognizes an epitope in the alpha subunits of CD221, demonstrated by western blotting (Li <i>et al.</i>1993). CD221 is expressed in a range of tissues, including kidney, liver, placenta, mammary gland, brain, ovary and skin. The ligands for CD221 include IGF-I and IGF-II, which bind to CD221 and activate tyrosine kinase activity, resulting in phosphorylation of several intracellular signaling proteins. Clone 1H7 is reported to partially block binding of IGF-I and IGF-II to CD221 (Li <i>et al.</i>1993).
Flow Cytometry	Use 10µl of the suggested working dilution to label 10^6 cells in 100µl
References	 Li, S.L. <i>et al.</i> (1993) Two new monoclonal antibodies against the alpha subunit of the human insulin-like growth factor-I receptor. <u>Biochem Biophys Res Commun. 196 (1): 92-8.</u> Beauvais, D.M. & Rapraeger, A.C. (2010) Syndecan-1 couples the insulin-like growth factor-1 receptor to inside-out integrin activation. <u>J Cell Sci. 123 (Pt 21): 3796-807.</u> Krieger, C.C. <i>et al.</i> (2016) Thyrotropin/IGF-1 receptor crosstalk in Graves' ophthalmopathy pathogenesis. <u>J Clin Endocrinol Metab. Apr 4: jc20161315. [Epub ahead of print]</u> van Valen, F. <i>et al.</i> (2012) A Novel Role of IGF1 in Apo2L/TRAIL-Mediated Apoptosis of Ewing Tumor Cells. <u>Sarcoma. 2012: 782970.</u>
Storage	Prior to reconstitution store at +4°C. Following reconstitution store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Health Ai Informati	-		•	sheet documentation antibodies.com/SDS/N	le at:
Regulato	ry	For resea	rch purpos	es only	
– – –	d Product	s			
Relate		0			
	nended Neg	-	ntrols		
Recomn		ative Co		<u> 4928PE)</u>	
Recomn	nended Ne		<u>_:RPE (MCA</u>	<u> 4928PE)</u>	
Recomn MOUSE Ig Recomn HUMAN S	nended Neg	ative Col CONTROI ful Reage	<u>_:RPE (MCA</u>	<u> 4928PE)</u>	

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint