

Datasheet: MCA2352GA

Description:	HAMSTER ANTI MOUSE CD121a
Specificity:	CD121a
Other names:	INTERLEUKIN 1 RECEPTOR TYPE 1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	JAMA-147
Isotype:	IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/25 - 1/100
Immunohistology - Frozen			•	
Immunohistology - Paraffin			•	
ELISA			•	
Immunoprecipitation	•			
Western Blotting			•	
Functional Assays (1)			•	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) Bio-Rad recommend the use of MCA2352XZ for functional assays.

Target Species	Mouse	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein G supernatant	From tissue culture
Buffer Solution	Phosphate buffered saline	
Preservative	0.09% sodium azide (NaN ₃)	

Stabilisers Approx. Protein IgG concentration 1.0 mg/ml Concentrations **Immunogen** Extracellular domain of murine IL-1R. **External Database UniProt:** Links P13504 Related reagents **Entrez Gene:** <u>16177</u> II1r1 Related reagents **Synonyms** II-1r1, II1ra **RRID** AB 566480 **Fusion Partners** Spleen cells from immunised Armenian Hamster were fused with cells of the mouse P3X63.Ag 8-653 myeloma cell line. **Specificity** Hamster anti Mouse CD121a monoclonal antibody, clone JAMA-147 recognizes murine CD121a, the ~80 kDa type I interleukin-1 receptor. CD121a is a single pass type I membrane glycoprotein, weakly expressed on most cells and tissues including T lymphocytes, thymocytes, epithelial cells and dendritic cells, with stronger expression reported on fibroblasts. The ligands for CD121a are the interleukin-1 (IL-1) family members; IL-1 alpha; IL-1 beta and IL-1 receptor antagonist. IL-1 is a mediator of inflammation and can elicit a variety of other biological responses through binding to CD121a. Hamster anti mouse CD121 antibody, clone JAMA-147 has been reported to block IL-1 dependent cellular responses (Rogers et al. 1992). Flow Cytometry Use 10µl of the suggested working dilution to label 10⁶ cells in 100µl References 1. Beckerman, K.P. et al. (1993) Release of nitric oxide during the T cell-independent pathway of macrophage activation. Its role in resistance to Listeria monocytogenes. J Immunol. 150 (3): 888-95. 2. Merrick, J.C. et al. (1997) Lymphocyte apoptosis during early phase of Listeria infection in mice. Am J Pathol. 151 (3): 785-92. 3. Scarim, A.L. et al. (1997) Evidence for the presence of type I IL-1 receptors on beta-cells of islets of Langerhans. Biochim Biophys Acta. 1361 (3): 313-20. 4. Kavanagh, D.P. et al. (2013) Mechanisms of adhesion and subsequent actions of a

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for

haematopoietic stem cell line, HPC-7, in the injured murine intestinal microcirculation in

vivo. PLoS One. 8 (3): e59150.

short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2352GA 10040
Regulatory	For research purposes only

Related Products

America

Recommended Secondary Antibodies

Goat Anti Hamster IgG (STAR104...) DyLight®550, DyLight®650, DyLight®800,

FITC

Goat Anti Hamster IgG (STAR79...) Biotin, FITC, HRP

Recommended Negative Controls

Fax: +1 919 878 3751

HAMSTER (ARMENIAN) IgG NEGATIVE CONTROL (MCA2356)

North & South Tel: +1 800 265 7376 Worldwide

Email: antibody_sales_us@bio-rad.com

Tel: +44 (0)1865 852 700 Europe Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M434200:250128'

Printed on 28 Jan 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint