

Datasheet: MCA2363A647

Description:	RAT ANTI MOUSE CD122:Alexa Fluor® 647
Specificity:	CD122
Other names:	INTERLEUKIN-2 RECEPTOR BETA CHAIN
Format:	ALEXA FLUOR® 647
Product Type:	Monoclonal Antibody
Clone:	TM-BETA 1
Isotype:	IgG2b
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system to a concentration equivalent to their test reagent.

Target Species	Mouse		
Product Form	Purified IgG conjugated to Alexa Fluor® 647- liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Alexa Fluor®647	650	665
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml		

Immunogen Cells from murine IL-2RB⁺ transformant TART-mBeta.

External Database

Links

UniProt:

[P16297](#)

[Related reagents](#)

Entrez Gene:

[16185](#)

Il2rb

[Related reagents](#)

RRID

AB_566483

Fusion Partners

Lymph node cells from immunised Sprague Dawley rats were fused with cells of the mouse PA1 myeloma cell line.

Specificity

Rat anti Mouse CD122 antibody, clone TM-BETA 1 recognizes the murine CD122 cell surface glycoprotein, a ~75-85 kDa molecule also known as interleukin-2 receptor (IL-2R) beta chain. In mouse, CD122 is primarily expressed on natural killer (NK) cells and at lower levels on lymphocyte subsets, monocytes and macrophages.

CD122 can associate with the IL-2R alpha chain (CD25) and CD132 to form a functional high affinity receptor complex that binds IL-2. CD132/CD122 heterodimers can also be formed which bind IL-2 with intermediate affinity. IL-2 binding to the IL-2R is important in the generation and differentiation of T lymphocytes and NK cells.

Rat anti Mouse CD122 antibody, clone TM-BETA 1 blocks IL-2 high affinity binding to the IL-2R beta receptor ([Tanaka *et al.* 1991](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR ([BUF041A/B](#)).

References

1. Tanaka, T. *et al.* (1991) A novel monoclonal antibody against murine IL-2 receptor beta-chain. Characterization of receptor expression in normal lymphoid cells and EL-4 cells. [J Immunol. 147 \(7\): 2222-8.](#)
2. Leung, K.T. (2011) The tetraspanin CD9 regulates migration, adhesion, and homing of human cord blood CD34+ hematopoietic stem and progenitor cells. [Blood. 117: 1840-50.](#)
3. Chenoweth, M.J. *et al.* (2012) IL-15 Can Signal via IL-15R α , JNK, and NF- κ B To Drive RANTES Production by Myeloid Cells. [J Immunol. 188: 4149-57.](#)
4. Mertens, J. *et al.* (2009) Streptococcus pneumoniae serotype 1 capsular polysaccharide induces CD8CD28 regulatory T lymphocytes by TCR crosslinking. [PLoS Pathog. 5 \(9\): e1000596.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in

frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Acknowledgements This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M385273:210513'

Printed on 21 Mar 2022

© 2022 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)