

## Datasheet: MCA2365PE

<b>Description:</b>	MOUSE ANTI BOVINE CD335:RPE
<b>Specificity:</b>	CD335
<b>Other names:</b>	NKp46
<b>Format:</b>	RPE
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	AKS1
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system to a concentration equivalent to their test reagent.

#### Target Species

Bovine

#### Species Cross Reactivity

Reacts with: American Bison

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

#### Product Form

Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized

#### Reconstitution

Reconstitute with 1 ml distilled water

#### Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
RPE 488nm laser	496	578

#### Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative</b>	0.09% sodium azide (NaN <sub>3</sub> )
<b>Stabilisers</b>	1% bovine serum albumin 5% sucrose
<b>Immunogen</b>	Fusion protein consisting of the extracellular region of bovine CD335.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">Q863H2</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">369024</a>    NCR1    <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_2149299
<b>Fusion Partners</b>	Spleen cells from immunised Balb/c mice were fused with cells of the NS-0 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Bovine CD335 antibody, clone AKS1</b> recognizes bovine CD335, also known as NKp46 and Natural cytotoxicity triggering receptor 1. CD335 is a type I transmembrane protein, with two extracellular C2-type immunoglobulin-like domains, which functions as an activating receptor. CD335 is expressed by human natural killer cells (<a href="#">Sivori <i>et al.</i> 1997</a>). The bovine homologue is expressed on bovine NK cells (<a href="#">Storset <i>et al.</i> 2004</a>) and no expression of CD335 has been detected on B cells, T cells, monocytes or granulocytes.</p> <p>Clone AKS1 is reported to activate lysis of FcγR-expressing cell line P815, by IL-2 activated NKp46+ cells (<a href="#">Storset <i>et al.</i> 2004</a>).</p>
<b>Flow Cytometry</b>	Use 10µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100µl
<b>References</b>	<ol style="list-style-type: none"> <li>1. Storset, A.K. <i>et al.</i> (2004) NKp46 defines a subset of bovine leukocytes with natural killer cell characteristics. <a href="#">Eur J Immunol. 34 (3): 669-76.</a></li> <li>2. Kulberg, S. <i>et al.</i> (2004) Reference values for relative numbers of natural killer cells in cattle blood. <a href="#">Dev Comp Immunol. 28 (9): 941-8.</a></li> <li>3. Van Rhijn, I. <i>et al.</i> (2007) Massive, sustained γδ T cell migration from the bovine skin <i>in vivo</i>. <a href="#">J Leukoc Biol. 81: 968-73.</a></li> <li>4. Toka, F.N. <i>et al.</i> (2011) Rapid and Transient Activation of γδ T Cells to IFN-γ Production, NK Cell-Like Killing, and Antigen Processing during Acute Virus Infection. <a href="#">J Immunol. 186: 4853-61.</a></li> <li>5. Bastos, R.G. <i>et al.</i> (2008) Bovine NK cells acquire cytotoxic activity and produce IFN-γ after stimulation by <i>Mycobacterium bovis</i> BCG- or <i>Babesia bovis</i>-exposed splenic dendritic cells. <a href="#">Vet Immunol Immunopathol. 124: 302-12.</a></li> <li>6. Boysen P <i>et al.</i> (2006) Bovine CD2-/NKp46+ cells are fully functional natural killer cells with a high activation status. <a href="#">BMC Immunol. 7: 10.</a></li> <li>7. Elh mouzi-Younes, J. <i>et al.</i> (2009) Bovine neonate natural killer cells are fully functional and highly responsive to interleukin-15 and to NKp46 receptor stimulation. <a href="#">Vet Res. 40:</a></li> </ol>

[54.](#)

8. Elh mouzi-Younes, J. *et al.* (2010) Ovine CD16+/CD14- blood lymphocytes present all the major characteristics of natural killer cells. [Vet Res. 41:4.](#)
9. Hoek, A. *et al.* (2009) Subpopulations of bovine WC1(+)  $\gamma/\delta$  T cells rather than CD4(+)CD25(high) Foxp3(+) T cells act as immune regulatory cells *ex vivo*. [Vet Res. 40:6.](#)
10. Zuerner, R.L. *et al.* (2011) A *Leptospira borgpetersenii* Serovar Hardjo Vaccine Induces a Th1 Response, Activates NK Cells, and Reduces Renal Colonization. [Clin Vaccine Immunol. 18: 684-91.](#)
11. Nelson, D.D. *et al.* (2010) CD8(+)/perforin(+)/WC1(-)  $\gamma/\delta$ T cells, not CD8(+)  $\alpha/\beta$  T cells, infiltrate vasculitis lesions of American bison (*Bison bison*) with experimental sheep-associated malignant catarrhal fever. [Vet Immunol Immunopathol. 136: 284-91.](#)
12. Banos, G. *et al.* (2013) Identification of immune traits correlated with dairy cow health, reproduction and productivity. [PLoS One. 8: e65766.](#)
13. Ramstead, A.G. *et al.* (2015) Aging influences the response of T cells to stimulation by the ellagitannin, oenothien B. [Int Immunopharmacol. 26 \(2\): 367-77.](#)
14. Boysen, P. *et al.* (2008) Natural killer cells in lymph nodes of healthy calves express CD16 and show both cytotoxic and cytokine-producing properties. [Dev Comp Immunol. 32 \(7\): 773-83.](#)
15. González-Cano, P. *et al.* (2014) Two functionally distinct myeloid dendritic cell subpopulations are present in bovine blood. [Dev Comp Immunol. 44 \(2\): 378-88.](#)
16. Di Palma, S. *et al.* (2012) Comparative spatiotemporal analysis of the intrathecal immune response in natural listeric rhombencephalitis of cattle and small ruminants. [Comp Immunol Microbiol Infect Dis. 35 \(5\): 429-41.](#)
17. Pirson, C. *et al.* (2015) Highly purified mycobacterial phosphatidylinositol mannosides drive cell-mediated responses and activate NKT cells in cattle. [Clin Vaccine Immunol. 22 \(2\): 178-84.](#)
18. Schneider, D.A. *et al.* (2011) Dynamics of bovine spleen cell populations during the acute response to *Babesia bovis* infection: an immunohistological study. [Parasite Immunol. 33 \(1\): 34-44.](#)
19. Van Meulder F *et al.* (2015) Analysis of the protective immune response following intramuscular vaccination of calves against the intestinal parasite *Cooperia oncophora*. [Int J Parasitol. 45 \(9-10\): 637-46.](#)
20. Patch, J.R. *et al.* (2014) Infection with foot-and-mouth disease virus (FMDV) induces a natural killer (NK) cell response in cattle that is lacking following vaccination. [Comp Immunol Microbiol Infect Dis. 37 \(4\): 249-57.](#)
21. Hedges, J.F. *et al.* (2015) Amphotericin B stimulates  $\gamma\delta$  T and NK cells, and enhances protection from *Salmonella* infection. [Innate Immun. 21 \(6\): 598-608.](#)
22. Allan, A.J. *et al.* (2015) Cattle NK Cell Heterogeneity and the Influence of MHC Class I. [J Immunol. 195 \(5\): 2199-206.](#)
23. Maślanka, T. *et al.* (2016) Prostaglandin E<sub>2</sub> exerts the proapoptotic and antiproliferative effects on bovine NK cells [Res Vet Sci. 107: 80-7.](#)
24. Hecker YP *et al.* (2015) Cell mediated immune responses in the placenta following challenge of vaccinated pregnant heifers with *Neospora caninum*. [Vet Parasitol. 214 \(3-4\): 247-54.](#)
25. Ohira K *et al.* (2016) Bovine leukemia virus reduces anti-viral cytokine activities and NK cytotoxicity by inducing TGF- $\beta$  secretion from regulatory T cells. [Immun Inflamm Dis. 4 \(1\): 52-63.](#)

26. Johnson, W.C. *et al.* (2008) Bovine WC1(-)  $\gamma\delta$ T cells incubated with IL-15 express the natural cytotoxicity receptor CD335 (NKp46) and produce IFN- $\gamma$  in response to exogenous IL-12 and IL-18. [Dev Comp Immunol. 32 \(8\): 1002-10.](#)
27. Sipka, A. *et al.* (2016) Bovine natural killer cells are present in *Escherichia coli* infected mammary gland tissue and show antimicrobial activity *in vitro*. [Comp Immunol, Microbiol Infect Dis. 48: 54-60.](#)
28. Hamilton, C.A. *et al.* (2016) Interactions between natural killer cells and dendritic cells favour T helper1-type responses to BCG in calves. [Vet Res. 47 \(1\): 85.](#)
29. Krueger, L.A. *et al.* (2016) Gamma delta T cells are early responders to *Mycobacterium avium* ssp. *paratuberculosis* in colostrum-replete Holstein calves. [J Dairy Sci. Sep 7. pii: S0022-0302\(16\)30611-7. \[Epub ahead of print\]](#)
30. Steinbach, S. *et al.* (2016) CD4+ and  $\gamma\delta$  T Cells are the main Producers of IL-22 and IL-17A in Lymphocytes from *Mycobacterium bovis*-infected Cattle. [Sci Rep. 6: 29990.](#)
31. Rutigliano, H.M. *et al.* (2016) Trophoblast Major Histocompatibility Complex Class I Expression Is Associated with Immune-Mediated Rejection of Bovine Fetuses Produced by Cloning. [Biol Reprod. 95 \(2\): 39.](#)
32. González-Hernández A *et al.* (2016) Host protective ASP-based vaccine against the parasitic nematode *Ostertagia ostertagi* triggers NK cell activation and mixed IgG1-IgG2 response. [Sci Rep. 6: 29496.](#)
33. Khosa, S. *et al.* (2020) Bovine Adenovirus-3 Tropism for Bovine Leukocyte Sub-Populations. [Viruses. 12 \(12\)Dec 12 \[Epub ahead of print\].](#)
34. Denholm, S.J. *et al.* (2017) Estimating genetic and phenotypic parameters of cellular immune-associated traits in dairy cows. [J Dairy Sci. 100 \(4\): 2850-2862.](#)
35. Hamilton, C.A. *et al.* (2017) Frequency and phenotype of natural killer cells and natural killer cell subsets in bovine lymphoid compartments and blood. [Immunology. 151 \(1\): 89-97.](#)
36. Bassi, P.B. *et al.* (2018) Parasitological and immunological evaluation of cattle experimentally infected with *Trypanosoma vivax*. [Exp Parasitol. 185: 98-106.](#)
37. de Araújo, F.F. *et al.* (2019) Distinct immune response profile during *Rhipicephalus* (*Boophilus*) *microplus*. infestations of guzerat dairy herd according to the maternal lineage ancestry (mitochondrial DNA). [Vet Parasitol. 273: 36-44.](#)
38. Nakajima, N. *et al.* (2019) Effects of direct exposure to cold weather under grazing in winter on the physiological, immunological, and behavioral conditions of Japanese Black beef cattle in central Japan. [Anim Sci J. 90 \(8\): 1033-1041.](#)
39. Okino, C.H. *et al.* (2020) A polymorphic CD4 epitope related to increased susceptibility to *Babesia bovis*. in Canchim calves. [Vet Immunol Immunopathol. 230: 110132.](#)
40. de Araújo, F.F. *et al.* (2019) Distinct immune response profile during *Rhipicephalus* (*Boophilus*) *microplus*. infestations of guzerat dairy herd according to the maternal lineage ancestry (mitochondrial DNA). [Vet Parasitol. 273: 36-44.](#)
41. Colombatti Olivieri, M.A. *et al.* (2021) Evaluation of a virulent strain of *Mycobacterium avium*. subsp. *paratuberculosis*. used as a heat-killed vaccine. [Vaccine. Nov 10;S0264-410X\(21\)01433-X.](#)

---

#### Further Reading

1. Sivori, S. *et al.* (1997) p46, a novel natural killer cell-specific surface molecule that mediates cell activation. [J Exp Med. 186 \(7\): 1129-36.](#)
2. Storset, A.K. *et al.* (2003) Natural killer cell receptors in cattle: a bovine killer cell immunoglobulin-like receptor multigene family contains members with divergent signaling

motifs. [Eur J Immunol. 33 \(4\): 980-90.](#)

---

**Storage** Prior to reconstitution store at +4°C.  
Following reconstitution store at +4°C.  
DO NOT FREEZE.  
This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #20487 available at:  
<https://www.bio-rad-antibodies.com/SDS/MCA2365PE>  
20487

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M414724:221213'

**Printed on 20 Apr 2023**

---

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)