

Datasheet: MCA2367F

Description:	RAT ANTI MOUSE CD197:FITC
Specificity:	CD197
Other names:	CCR7
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	4B12
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

RRID AB_566602

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat *

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species Mouse

Product Form Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

Preparation Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution Phosphate buffered saline

Preservative 0.09% Sodium Azide
Stabilisers 1% Bovine Serum Albumin

Approx. Protein Concentrations IgG concentration 0.1 mg/ml

Immunogen CCR7 rat basophilic leukaemia (RBL)-2H3 transfectants.

External Database Links **UniProt:**

[P47774](#) [Related reagents](#)

Entrez Gene:

[12775](#) Ccr7 [Related reagents](#)

Synonyms	Cmkbr7, Ebi1, Ebi1h
-----------------	---------------------

Fusion Partners	Spleen cells from immunised Lou/c rats were fused with cells of the mouse X63.Ag8.653 myeloma cell line.
------------------------	----------------------------------------------------------------------------------------------------------

Specificity	<p>Rat anti Mouse CD197 antibody, clone 4B12 recognizes murine C-C chemokine receptor type 7, also known as CD197, Epstein-Barr virus-induced G-protein coupled receptor 1, CCR7 or MIP-3 beta receptor. CD197 is a 354 amino acid ~43 kDa member of the G-protein coupled receptor family. In mice, CD197 is expressed on most naïve T cells, a subset of memory T cells, B cells, mature dendritic cells, NK cells, a subset of thymocytes and CD34+ macrophage progenitor cells.</p> <p>CD197 is a functional receptor for chemokines CCL21 and CCL19. The receptor plays an important role in regulating the trafficking and homing of lymphocytes and dendritic cells to secondary lymphoid organs.</p> <p>Rat anti Mouse CD197 antibody, clone 4B12 does not block the binding of the ligand to CD197.</p>
--------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Flow Cytometry	<p>Use 20ul of the suggested working dilution to label 10⁶ cells in 100ul.</p> <p>The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B).</p>
-----------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

References	1. Ritter, U. <i>et al.</i> (2004) Analysis of the CCR7 expression on murine bone marrow-derived and spleen dendritic cells. J Leukoc Biol. 76 (2): 472-6.
-------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
----------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Guarantee	18 months from date of despatch.
------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
--------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:FITC \(MCA1212F\)](#)

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M342161:190110'

Printed on 20 May 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)