

Datasheet: MCA2367PE

Description:	RAT ANTI MOUSE CD197:RPE			
Specificity:	CD197			
Other names:	CCR7			
Format:	RPE			
Product Type:	Monoclonal Antibody			
Clone:	4B12			
Isotype:	lgG2a			
Quantity:	100 TESTS			

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .							
		Yes	No	Not Determined	Suggested Dilution			
	Flow Cytometry	-			Neat			
	Where this product ha			•	•			
	necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.							
Target Species	Mouse							
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized							
Reconstitution	Reconstitute with 1.0 ml distilled water							
Max Ex/Em	Fluorophore	Excitation Ma	ax (nm)	Emission Max (nm)				
	RPE 488nm laser	496		578				
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant							
Buffer Solution	Phosphate buffered saline							
Preservative Stabilisers	0.09% sodium azide (NaN ₃) 1% bovine serum albumin 5% sucrose							

External Database					
Links	UniProt: P47774 Related reagents				
	Entrez Gene:				
	<u>12775</u> Ccr7 <u>Related reagents</u>				
Synonyms	Cmkbr7, Ebi1, Ebi1h				
RRID	AB_566600				
Fusion Partners	Spleen cells from immunised Lou/c rats were fused with cells of the mouse X63.Ag8.653 myeloma cell line.				
Specificity	Rat anti Mouse CD197 antibody, clone 4B12 recognizes murine C-C chemokine receptor type 7, also known as CD197, Epstein-Barr virus-induced G-protein coupled receptor 1, CCR7 or MIP-3 beta receptor. CD197 is a 354 amino acid ~43 kDa member of the G-protein coupled receptor family. In mice, CD197 is expressed on most naïve T cells, a subset of memory T cells, B cells, mature dendritic cells, NK cells, a subset of thymocytes and CD34+ macrophage progenitor cells.				
	important role in regulating the trafficking and homing of lymphocytes and dendritic cells to secondary lymphoid organs. Rat anti Mouse CD197 antibody, clone 4B12 does not block the binding of the ligand to CD197.				
Flow Cytometry	Use 20µl of the suggested working dilution to label 10 ⁶ cells in 100µl. The Fc region of monoclonal antibodies may bind to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (<u>BUF041A/BUF041B</u>).				
References	1. Ritter, U. <i>et al.</i> (2004) Analysis of the CCR7 expression on murine bone marrow- derived and spleen dendritic cells. <u>J Leukoc Biol. 76 (2): 472-6.</u>				
Storage	Store at +4°C. DO NOT FREEZE This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.				
Guarantee	12 months from date of despatch				
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA2367PE 20487				

Related Products

Recommended Negative Controls

RAT IgG2a NEGATIVE CONTROL:RPE (MCA1212PE)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M419425:230616'

Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint