

Datasheet: MCA2367PET

Description:	RAT ANTI MOUSE CD197:RPE	
Specificity:	CD197	
Other names:	CCR7	
Format:	RPE	
Product Type:	Monoclonal Antibody	
Clone:	4B12	
Isotype:	lgG2a	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

External Database Links

UniProt:

P47774 Related reagents

Entrez Gene:

12775 Ccr7 Related reagents

Synonyms	Cmkbr7, Ebi1, Ebi1h
Fusion Partners	Spleen cells from immunised Lou/c rats were fused with cells of the mouse X63.Ag8.653 myeloma cell line.
Specificity	Rat anti Mouse CD197 antibody, clone 4B12 recognizes murine C-C chemokine receptor type 7, also known as CD197, Epstein-Barr virus-induced G-protein coupled receptor 1, CCR7 or MIP-3 beta receptor. CD197 is a 354 amino acid ~43 kDa member of the G-protein coupled receptor family. In mice, CD197 is expressed on most naïve T cells, a subset of memory T cells, B cells, mature dendritic cells, NK cells, a subset of thymocytes and CD34+ macrophage progenitor cells.
	CD197 is a functional receptor for chemokines CCL21 and CCL19. The receptor plays an important role in regulating the trafficking and homing of lymphocytes and dendritic cells to secondary lymphoid organs.
	Rat anti Mouse CD197 antibody, clone 4B12 does not block the binding of the ligand to CD197.
Flow Cytometry	Use 20ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
	The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (<u>BUF041A/B</u>).
References	1. Ritter, U. <i>et al.</i> (2004) Analysis of the CCR7 expression on murine bone marrow-derived and spleen dendritic cells. <u>J Leukoc Biol. 76 (2): 472-6.</u>
Storage	Store at +4°C.
	DO NOT FREEZE
	This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of reconstitution.
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
Regulatory	

Related Products

Recommended Negative Controls

RAT IgG2a NEGATIVE CONTROL:RPE (MCA1212PE)

Recommended Useful Reagents

MOUSE SEROBLOCK FcR (BUF041A)
MOUSE SEROBLOCK FcR (BUF041B)

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 **Europe** Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_uk@bio-rad.com

'M323728:180727'

Email: antibody_sales_de@bio-rad.com

Printed on 06 Dec 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint

Email: antibody_sales_us@bio-rad.com