

Datasheet: MCA2368PET

Description:	MOUSE ANTI HUMAN FC EPSILON R1 ALPHA:RPE		
Specificity:	Fc EPSILON R1 ALPHA		
Format:	RPE		
Product Type:	Monoclonal Antibody		
Clone:	9E1		
Isotype:	lgG2b		
Quantity:	25 TESTS		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	=			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species						
Product Form	Purified IgG conjug	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized				
Reconstitution	Reconstitute in 0.25 ml disilled water					
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)			
	RPE 488nm laser	496	578			
Preparation	on Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatar					
Buffer Solution	Phosphate buffered	saline				
Preservative	0.09% Sodium Azio	le				
Stabilisers	1% Bovine Seru					
	5% Sucrose					
External Database Links	UniProt:					

2205 FCER1A Related reagents

Related reagents

P12319

Entrez Gene:

Synonyms	FCE1A
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the Sp2/0-Ag14 myeloma cell line.
Specificity	Mouse anti Human Fc epsilon R1 alpha antibody, clone 9E1 recognizes the human high affinity Fc receptor for IgE (Fc epsilon R1), which plays a central role in the IgE-mediated allergic response. Fc epsilon R1 consists of four subunits, a high affinity IgE binding alpha subunit, a beta chain and two di-sulphide linked gamma subunits. Clone 9E1 specifically recognizes the extracellular D1 domain of the Fc epsilon R1 alpha chain.
	Fc epsilon R1 is primarily expressed on mast cells and basophils but expression of Fc epsilon R1 has also been reported on monocytes, Langerhans cells and dendritic cells from patients with atopic diseases.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	 Vangelista, L. et al. (2002) A minimal receptor-Ig chimera of human FcepsilonRI alpha-chain efficiently binds secretory and membrane IgE. Protein Eng. 15 (1): 51-7. Vangelista, L. et al. (2002) Efficient folding of the FcepsilonRI alpha-chain membrane-proximal domain D2 depends on the presence of the N-terminal domain D1. J Mol Biol. 322 (4): 815-25. Vangelista, L. et al. (2005) Membrane IgE binds and activates Fc epsilon RI in an antigen-independent manner. J Immunol. 174 (9): 5602-11.
Storage	Prior to reconstitution store at +4°C. After reconstitution store at +4°C. DO NOT FREEZE. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of reconstitution.
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG2b NEGATIVE CONTROL:RPE (MCA691PE)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376 America

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

'M323732:180727'

Printed on 06 Dec 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint