

Datasheet: MCA2372B

## **BATCH NUMBER 1804**

Description:	MOUSE ANTI BOVINE INTERLEUKIN-4:Biotin	
Specificity:	IL-4	
Format:	Biotin	
<b>Product Type:</b>	Monoclonal Antibody	
Clone:	CC314	
Isotype:	lgG2b	
Quantity:	0.25 mg	

## **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen			•	
Immunohistology - Paraffin			•	
ELISA	•			1ug/ml - 5ug/ml
Immunoprecipitation			•	
Western Blotting			•	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Bovine
Species Cross Reactivity	Reacts with: Goat, Sheep <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG conjugated to Biotin - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Recombinant bovine IL-4.
External Database Links	UniProt: P30367 Related reagents  Entrez Gene: 280824 IL4 Related reagents
RRID	AB_2127032
Fusion Partners	Spleen cells form immunised BALB/c mice were fused with cells of the SP2/0 myeloma cell line.
Specificity	Mouse anti Bovine interleukin-4 antibody, clone CC314 recognizes bovine intetleukin-4 (IL-4), also known as B-cell stimulatory factor 1. IL-4 is expressed by a range of cells including T and B cells, macrophages and monocytes. It plays an important role in the regulation of bovine T and B cell responses. Bovine interleukin-4 is a 15 kDa, 135 amino acid peptide which is processed to a mature secreted form with the cleavage of a 24 amino acid, N-terminal signal peptide resulting in a 12.6 kDa active cytokine, the apparent MW may be altered by glycosylation (Heussler et al. 1992).
	IL-4 is expressed by a range of cells including T and B cells, macrophages and monocytes. It plays an important role in the regulation of bovine T and B cell responses.
	Mouse anti Bovine interleukin-4, clone CC314 has been used successfully as a detection antibody in a sandwich ELISA with Mouse anti Bovine IL-4 antibody, clone CC313 (MCA2371) as a coating reagent both with bovine (Stabel et al. 2011) and ovine samples (Olivier et al. 2012). Clones CC313 and CC314 have been used in the reverse configuration for the measurement of interleukin-4 in caprine samples (Marinaro et al. 2012).
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.
ELISA	This biotin conjugate may be used as detection reagent in sandwich ELISA assays for bovine IL-4 with MCA2371 as capture reagent, and recombinant bovine IL-4 (PBP006) as a standard.
References	1. Hope, J.C. <i>et al.</i> (2005) Development of detection methods for ruminant interleukin (IL)-4. <u>J Immunol Methods</u> . 301 (1-2): 114-23.

- 2. Abbott JR et al. (2005) Rapid and long-term disappearance of CD4+ T lymphocyte responses specific for Anaplasma marginale major surface protein-2 (MSP2) in MSP2 vaccinates following challenge with live A. marginale. J Immunol. 174 (11): 6702-15.
- 3. Stabel JR et al. (2013) Disparate Host Immunity to Mycobacterium avium subsp. paratuberculosis Antigens in Calves Inoculated with M. avium subsp. paratuberculosis, M. avium subsp. avium, M. kansasii and M. bovis. Clin Vaccine Immunol. 20 (6): 848-57.
- 4. Stabel, J.R. et al. (2011) Mediation of host immune responses after immunization of neonatal calves with a heat-killed Mycobacterium avium subsp. paratuberculosis vaccine. Clin Vaccine Immunol. 18: 2079-89.
- 5. Marinaro, M. et al. (2012) Antigen-specific IFN-gamma and IL-4 production in caprine herpesvirus infected goats. Res Vet Sci. 93: 662-7.
- 6. Olivier, M. et al. (2012) Capacities of migrating CD1b+ lymph dendritic cells to present Salmonella antigens to naive T cells. PLoS One. 7: e30430.
- 7. Cassady-Cain, R.L. et al. (2017) Inhibition of Antigen-Specific and Nonspecific Stimulation of Bovine T and B Cells by Lymphostatin from Attaching and Effacing Escherichia coli. Infect Immun. 85 (2): pii: e00845-16. [Epub ahead of print]

Storage	Store at +4°C or at -20°C if preferred.
	Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2372B">https://www.bio-rad-antibodies.com/SDS/MCA2372B</a> 10041
Regulatory	For research purposes only

North & South Tel: +1 800 265 7376 Tel: +44 (0)1865 852 700 Worldwide Europe Tel: +49 (0) 89 8090 95 21 America Fax: +49 (0) 89 8090 95 50 Fax: +1 919 878 3751 Fax: +44 (0)1865 852 739

> Email: antibody\_sales\_us@bio-rad.com Email: antibody\_sales\_uk@bio-rad.com Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M366736:200529'

## Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint