

# Datasheet: MCA2385PE BATCH NUMBER 164046

MOUSE ANTI HORSE CD8:RPE
CD8
RPE
Monoclonal Antibody
CVS8
lgG1
100 TESTS

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further							
	information. For general protocol recommendations, please visit <u>www.bio-</u> rad-antibodies.com/protocols.							
		Yes	No	Not Determined	Suggested Dilution			
	Flow Cytometry	•			Neat			
	Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.							
Target Species	Horse							
Species Cross	Reacts with: Ass Does not react with:Zebra							
Reactivity								
	N.B. Antibody reactivi				•			
	reactivity is derived fro	•		· •	•			
	personal communications from the originators. Please refer to references indicated for further information.							
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - Iyophilized							
Reconstitution	Reconstitute with 1.0 ml distilled water							
	Care should be taken during reconstitution as the protein may appear as a film at the							
	bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution							
Max Ex/Em	Fluorophore	Excitation Ma	x (nm)	Emission Max (nm)				
	RPE 488nm laser	496		578				

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> ) 1% Bovine Serum Albumin 5% Sucrose
Immunogen	Equine PBMCs.
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the X63-Ag8.653 myeloma cell line.
Specificity	<b>Mouse anti Horse CD8 (clone CVS8),</b> is a monoclonal antibody recognising the equine homologue of the human CD8 cell surface antigen which is expressed by a subset of T lymphocytes.
	A study undertaken using CVS8 to identify CD8 on several wild african equid species indicates that the CVS8 clone recognizes Somali wild ass ( <i>Equus asinus</i> ) but not Grévy's Zebra ( <i>E. grevyi</i> ) or Hartmann's Mountain Zebra ( <i>E. zebra</i> ) ( <u>Ibrahim 2007</u> ).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
References	<ol> <li>Lunn, D.P et al (1991) Three monoclonal antibodies identifying antigens on all equine T lymphocytes, and two mutually exclusive T-lymphocyte subsets <u>Immunology 74: 251-257.</u></li> <li>Jacks, S. (2007) Experimental infection of neonatal foals with Rhodococcus equi triggers adult-like gamma interferon induction. <u>Clin Vaccine Immunol. 14: 669-77.</u></li> <li>Pearson, W. <i>et al.</i> (2007) Low-dose ginseng (Panax quinquefolium) modulates the course and magnitude of the antibody response to vaccination against equid herpesvirus I in horses. <u>Can J Vet Res. 71: 213-7.</u></li> <li>Lunn, D.P. <i>et al.</i> (1998) Report of the second equine leucocyte antigen workshop, Squaw Valley, California July 1995. <u>Vet Immunol Immunopathol. 62: 101-143.</u></li> <li>Merant, C. <i>et al.</i> (2003) Cross-species reactivity of seven monoclonal antibodies with equine lymphocytes by flow cytometry. <u>Vet Res. 34: 791-801.</u></li> <li>Ibrahim, S (2007) Analysis of monoclonal antibody cross-reactivity with leukocytes from equids and cloning of CD28 <u>Chapter 5 in PhD Thesis Freie Universität Berlin</u></li> <li>Carossino, M. <i>et al.</i> (2019) Equine arteritis virus long-term persistence is orchestrated by CD8+ T lymphocyte transcription factors, inhibitory receptors, and the CXCL16/CXCR6 axis. <u>PLoS Pathog. 15 (7): e1007950.</u></li> <li>Tomlinson, J.E. <i>et al.</i> (2018) Multispectral fluorescence-activated cell sorting of B and T cell subpopulations from equine peripheral blood. <u>Vet Immunol Immunopathol. 199: 22-31.</u></li> <li>Placci, M. <i>et al.</i> (2020) Natural Horse Boarding Vs Traditional Stable: A Comparison of Hormonal, Hematological and Immunological Parameters. <u>J Appl Anim Welf Sci. 23 (3):</u> <u>366-77.</u></li> </ol>
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be

This product should be stored undiluted. This product is photosensitive and should be

	protected from light.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA2385PE 20487
Regulatory	For research purposes only

### **Related Products**

### **Recommended Useful Reagents**

### MOUSE ANTI HORSE CD4:FITC (MCA1078F)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-r	ad.com	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M375471:210104'

#### Printed on 05 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint