

Datasheet: MCA2385PE

BATCH NUMBER INN1704

Description:	MOUSE ANTI HORSE CD8:RPE
Specificity:	CD8
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	CVS8
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Horse		
Species Cross Reactivity	Reacts with: Ass Does not react with:Zebra N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1.0 ml distilled water Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin 5% Sucrose
Immunogen	Equine PBMCs.
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the X63-Ag8.653 myeloma cell line.
Specificity	<p>Mouse anti Horse CD8 (clone CVS8), is a monoclonal antibody recognising the equine homologue of the human CD8 cell surface antigen which is expressed by a subset of T lymphocytes.</p> <p>A study undertaken using CVS8 to identify CD8 on several wild african equid species indicates that the CVS8 clone recognizes Somali wild ass (<i>Equus asinus</i>) but not Grévy's Zebra (<i>E. grevyi</i>) or Hartmann's Mountain Zebra (<i>E. zebra</i>) (Ibrahim 2007).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Lunn, D.P et al (1991) Three monoclonal antibodies identifying antigens on all equine T lymphocytes, and two mutually exclusive T-lymphocyte subsets Immunology 74: 251-257. 2. Jacks, S. (2007) Experimental infection of neonatal foals with Rhodococcus equi triggers adult-like gamma interferon induction. Clin Vaccine Immunol. 14: 669-77. 3. Pearson, W. <i>et al.</i> (2007) Low-dose ginseng (Panax quinquefolium) modulates the course and magnitude of the antibody response to vaccination against equid herpesvirus 1 in horses. Can J Vet Res. 71: 213-7. 4. Lunn, D.P. <i>et al.</i> (1998) Report of the second equine leucocyte antigen workshop, Squaw Valley, California July 1995. Vet Immunol Immunopathol. 62: 101-143. 5. Merant, C. <i>et al.</i> (2003) Cross-species reactivity of seven monoclonal antibodies with equine lymphocytes by flow cytometry. Vet Res. 34: 791-801. 6. Ibrahim, S (2007) Analysis of monoclonal antibody cross-reactivity with leukocytes from equids and cloning of CD28 Chapter 5 in PhD Thesis Freie Universität Berlin 7. Carossino, M. <i>et al.</i> (2019) Equine arteritis virus long-term persistence is orchestrated by CD8+ T lymphocyte transcription factors, inhibitory receptors, and the CXCL16/CXCR6 axis. PLoS Pathog. 15 (7): e1007950. 8. Tomlinson, J.E. <i>et al.</i> (2018) Multispectral fluorescence-activated cell sorting of B and T cell subpopulations from equine peripheral blood. Vet Immunol Immunopathol. 199: 22-31. 9. Placci, M. <i>et al.</i> (2020) Natural Horse Boarding Vs Traditional Stable: A Comparison of Hormonal, Hematological and Immunological Parameters. J Appl Anim Welf Sci. 23 (3): 366-77.
Storage	<p>Store at +4°C. DO NOT FREEZE.</p> <p>This product should be stored undiluted. This product is photosensitive and should be</p>

protected from light.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA2385PE20487
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[MOUSE ANTI HORSE CD4:FITC \(MCA1078F\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M375471:210104'

Printed on 05 Feb 2024