

Datasheet: MCA2385PE

Description:	MOUSE ANTI HORSE CD8:RPE
Specificity:	CD8
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	CVS8
Isotype:	lgG1
Quantity:	100 TESTS

#### **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Horse						
Species Cross	Reacts with: Ass						
Reactivity	Does not react with:Zebra						
	reactivity is derived	from testing within our la	ons may vary between species. Cros aboratories, peer-reviewed publication rs. Please refer to references indicate	ns or			
Product Form	Purified IgG conjuga	ated to R. Phycoerythrin	(RPE) - lyophilized				
Reconstitution	Reconstitute with 1.	.0 ml distilled water					
	Care should be take	en during reconstitution	as the protein may appear as a film a	t the			
	bottom of the vial. E	Bio-Rad recommend that	the vial is gently mixed after reconsti	itution.			
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)				
	RPE 488nm laser	496	578				

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	<ul><li>0.09% Sodium Azide (NaN<sub>3</sub>)</li><li>1% Bovine Serum Albumin</li><li>5% Sucrose</li></ul>
Immunogen	Equine PBMCs.
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the X63-Ag8.653 myeloma cell line.
Specificity	<b>Mouse anti Horse CD8 (clone CVS8),</b> is a monoclonal antibody recognising the equine homologue of the human CD8 cell surface antigen which is expressed by a subset of T lymphocytes.
	A study undertaken using CVS8 to identify CD8 on several wild african equid species indicates that the CVS8 clone recognizes Somali wild ass ( <i>Equus asinus</i> ) but not Grévy's Zebra ( <i>E. grevyi</i> ) or Hartmann's Mountain Zebra ( <i>E. zebra</i> ) ( <u>Ibrahim 2007</u> ).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
References	1. Lunn, D.P et al (1991) Three monoclonal antibodies identifying antigens on all equine T lymphocytes, and two mutually exclusive T-lymphocyte subsets Immunology 74: 251-257.  2. Jacks, S. (2007) Experimental infection of neonatal foals with Rhodococcus equi triggers adult-like gamma interferon induction. Clin Vaccine Immunol. 14: 669-77.  3. Pearson, W. et al. (2007) Low-dose ginseng (Panax quinquefolium) modulates the course and magnitude of the antibody response to vaccination against equid herpesvirus I in horses. Can J Vet Res. 71: 213-7.  4. Lunn, D.P. et al. (1998) Report of the second equine leucocyte antigen workshop, Squaw Valley, California July 1995. Vet Immunol Immunopathol. 62: 101-143.  5. Merant, C. et al. (2003) Cross-species reactivity of seven monoclonal antibodies with equine lymphocytes by flow cytometry. Vet Res. 34: 791-801.  6. Ibrahim, S (2007) Analysis of monoclonal antibody cross-reactivity with leukocytes from equids and cloning of CD28 Chapter 5 in PhD Thesis Freie Universität Berlin  7. Carossino, M. et al. (2019) Equine arteritis virus long-term persistence is orchestrated by CD8+ T lymphocyte transcription factors, inhibitory receptors, and the CXCL16/CXCR6 axis. PLoS Pathog. 15 (7): e1007950.  8. Tomlinson, J.E. et al. (2018) Multispectral fluorescence-activated cell sorting of B and T cell subpopulations from equine peripheral blood. Vet Immunol Immunopathol. 199: 22-31.  9. Placci, M. et al. (2020) Natural Horse Boarding Vs Traditional Stable: A Comparison of Hormonal, Hematological and Immunological Parameters. J Appl Anim Welf Sci. 23 (3): 366-77.

# Storage

Store at +4°C. DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be

#### protected from light.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: 20487: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/20487.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/20487.pdf</a>
Regulatory	For research purposes only

## **Related Products**

## **Recommended Useful Reagents**

## MOUSE ANTI HORSE CD4:FITC (MCA1078F)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M375471:210104'

#### Printed on 07 Jan 2022

© 2022 Bio-Rad Laboratories Inc | Legal | Imprint