

Datasheet: MCA2386EL

| | |
|----------------------|------------------------------------|
| Description: | RAT ANTI MOUSE CD223:Low Endotoxin |
| Specificity: | CD223 |
| Other names: | LAG-3 |
| Format: | Low Endotoxin |
| Product Type: | Monoclonal Antibody |
| Clone: | C9B7W |
| Isotype: | IgG1 |
| Quantity: | 0.5 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | 1/50 - 1/200 |
| Immunohistology - Frozen | | | ▪ | |
| Immunohistology - Paraffin | | | ▪ | |
| ELISA | | | ▪ | |
| Immunoprecipitation | | | ▪ | |
| Western Blotting | | | ▪ | |
| Functional Assays | ▪ | | | |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| | |
|---------------------------------|---|
| Target Species | Mouse |
| Product Form | Purified IgG - liquid |
| Preparation | Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant |
| Buffer Solution | Phosphate buffered saline |
| Preservative Stabilisers | None present |

| | |
|---------------------------------------|---|
| Carrier Free | Yes |
| Endotoxin Level | < 0.01 EU/μg |
| Approx. Protein Concentrations | IgG concentration 1.0 mg/ml |
| Immunogen | Murine CD223 Ig fusion protein. |
| External Database Links | <p>UniProt: Q61790 Related reagents</p> <p>Entrez Gene: 16768 Lag3 Related reagents</p> |
| RRID | AB_566654 |
| Fusion Partners | Cells from immunised Lewis rats were fused with cells of the Sp/20 myeloma cell line. |
| Specificity | <p>Rat anti Mouse CD223 antibody, clone C9B7W recognizes murine lymphocyte activation gene-3 (LAG-3), a ~70 kDa activation-induced cell surface molecule that is also referred to as CD223.</p> <p>Murine CD223 is expressed on activated CD4 positive and CD8 positive alpha/beta T lymphocytes and a subset of natural killer (NK) cells. CD223 binds to MHC class II molecules with high affinity and is reported to negatively regulate T cell homeostasis and T cell expansion.</p> <p>Clone C9B7W recognizes an epitope within the D2 domain of CD223. Rat anti Mouse CD223 antibody, clone C9B7W is reported to block the <i>in vitro</i> function of murine LAG-3 but does not block binding of LAG-3 to MHC class II (Workman et al. 2002).</p> |
| Flow Cytometry | Use 10μl of the suggested working dilution to label 10 ⁶ cells in 100μl |
| References | <ol style="list-style-type: none"> 1. Workman, C.J. <i>et al.</i> (2002) Cutting edge: molecular analysis of the negative regulatory function of lymphocyte activation gene-3. J Immunol. 169 (10): 5392-5. 2. Workman, C.J. & Vignali, D.A. (2005) Negative regulation of T cell homeostasis by lymphocyte activation gene-3 (CD223). J Immunol. 174 (2): 688-95. 3. Byrne, K.T. <i>et al.</i> (2011) Autoimmune melanocyte destruction is required for robust CD8+ memory T cell responses to mouse melanoma. J Clin Invest. 121 (5): 1797-809. 4. Ordway, D. <i>et al.</i> (2007) The hypervirulent <i>Mycobacterium tuberculosis</i> strain HN878 induces a potent TH1 response followed by rapid down-regulation. J Immunol. 179: 522-31. 5. Hu, Z. <i>et al.</i> (2013) Regulatory CD8+ T cells associated with erosion of immune surveillance in persistent virus infection suppress <i>in vitro</i> and have a reversible proliferative defect. J Immunol. 191 (1): 312-22. 6. Iwasaki, Y. <i>et al.</i> (2013) Egr-2 transcription factor is required for Blimp-1-mediated IL-10 |

production in IL-27-stimulated CD4+ T cells. [Eur J Immunol. 43: 1063-73.](#)
7. Woo, S.R. *et al.* (2010) Differential subcellular localization of the regulatory T-cell protein LAG-3 and the coreceptor CD4. [Eur J Immunol. 40: 1768-77.](#)

| | |
|--------------------------------------|--|
| Storage | Store at -20°C only. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. |
| Guarantee | 12 months from date of despatch |
| Health And Safety Information | Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/MCA2386EL 10162 |
| Regulatory | For research purposes only |

Related Products

Recommended Secondary Antibodies

| | |
|--|---|
| Goat Anti Rat IgG (STAR73...) | RPE |
| Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) | DyLight®550 , DyLight®650 , DyLight®800 |
| Rabbit Anti Rat IgG (STAR21...) | HRP |
| Rabbit Anti Rat IgG (STAR16...) | DyLight®800 |
| Goat Anti Rat IgG (STAR131...) | Alk. Phos. , Biotin |
| Rabbit Anti Rat IgG (STAR17...) | FITC |
| Goat Anti Rat IgG (STAR72...) | HRP |
| Goat Anti Rat IgG (STAR69...) | FITC |

| | | | | | |
|----------------------------------|---|------------------|---|---------------|---|
| North & South America | Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com | Worldwide | Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com | Europe | Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com |
|----------------------------------|---|------------------|---|---------------|---|

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M414816:221213'

Printed on 04 Sep 2023