

Datasheet: MCA2387T

Description:	RAT ANTI MOUSE Gr-1
Specificity:	Gr-1
Other names:	Ly-6G
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	RB6-8C5
Isotype:	IgG2b
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/100 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃)

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Normal murine bone marrow cells.
External Database Links	<p>UniProt: P35461 Related reagents</p> <p>Entrez Gene: 546644 Ly6g Related reagents</p>
RRID	AB_2115659
Specificity	<p>Rat anti Mouse Gr-1 antibody, clone RB6-8C5 recognizes the mouse Gr-1 antigen, a ~21–25 kDa GPI anchored cell surface protein bearing a single uPAR/Ly6 domain that belongs to the Ly-6 family of proteins (Lee et al. 2013). Rat anti Mouse Gr-1 antibody, clone RB6-8C5 reacts predominantly with the Ly-6G protein but weaker reactivity with the Ly-6C protein has been reported (Fleming et al. 1993). However, other observations dispute the cross-reactivity of clone RB6-8C5 with the Ly-6C protein with the alternative explanation that certain sub-populations of bone marrow cells simultaneously express both Ly-6C and Ly-6G (Nagendra et al. 2007)</p> <p>The Gr-1 antigen is primarily a marker of myeloid differentiation. In the bone marrow the level of Gr-1 expression is low on immature myeloblasts and increases as the myeloid cells mature to granulocytes. Gr-1 is also expressed on macrophages and transiently on differentiating monocytes.</p> <p>Rat anti Mouse Gr-1 antibody, clone RB6-8C5 has been used successfully for the depletion of mature neutrophils <i>in vivo</i> (Czuprynski et al 1994, Daley et al. 2008).</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> Fleming, T.J. <i>et al.</i> (1993) Selective expression of Ly-6G on myeloid lineage cells in mouse bone marrow. RB6-8C5 mAb to granulocyte-differentiation antigen (Gr-1) detects members of the Ly-6 family. J Immunol. 151 (5): 2399-408. Hestdal, K. <i>et al.</i> (1991) Characterization and regulation of RB6-8C5 antigen expression on murine bone marrow cells. J Immunol. 147 (1): 22-8. Czuprynski, C.J. <i>et al.</i> (1994) Administration of anti-granulocyte mAb RB6-8C5 impairs the resistance of mice to <i>Listeria monocytogenes</i> infection. J Immunol. 152 (4): 1836-46. Sumagin R <i>et al.</i> (2010) LFA-1 and Mac-1 define characteristically different intraluminal crawling and emigration patterns for monocytes and neutrophils <i>in situ</i>. J Immunol. 185 (11): 7057-66. Takano, K. <i>et al.</i> (2011) Successful treatment of acute lung injury with pitavastatin in septic mice: potential role of glucocorticoid receptor expression in alveolar macrophages. J Pharmacol Exp Ther. 336: 381-90.

6. Giroux, M. *et al.* (2011) SMAD3 prevents graft-versus-host disease by restraining Th1 differentiation and granulocyte-mediated tissue damage. [Blood.117: 1734-44.](#)
7. Suttman, H. *et al.* (2006) Neutrophil granulocytes are required for effective Bacillus Calmette-Guérin immunotherapy of bladder cancer and orchestrate local immune responses. [Cancer Res. 66: 8250-7.](#)
8. Nix, R.N. *et al.* (2007) Hemophagocytic macrophages harbor *Salmonella enterica* during persistent infection. [PLoS Pathog. 3: e193.](#)
9. Kanda, N. *et al.* (2011) Visfatin Enhances CXCL8, CXCL10, and CCL20 Production in Human Keratinocytes. [Endocrinology. 152: 3155-64.](#)
10. Conlan, J. and North, R. (1994) Neutrophils are essential for early anti-*Listeria* defense in the liver, but not in the spleen or peritoneal cavity, as revealed by a granulocyte-depleting monoclonal antibody. [J Exp Med. 179:259-68.](#)
11. Takebe, M. *et al.* (2014) Inhibition of histone deacetylases protects septic mice from lung and splenic apoptosis. [J Surg Res. 187 \(2\): 559-70.](#)
12. Francke, A. *et al.* (2011) Generation of mature murine monocytes from heterogeneous bone marrow and description of their properties. [J Histochem Cytochem. 59: 813-25.](#)
13. Sharp, P.E. *et al.* (2013) FcγRIIb on myeloid cells and intrinsic renal cells rather than B cells protects from nephrotoxic nephritis. [J Immunol.190: 340-8.](#)
14. Hamers, A.A. *et al.* (2014) Limited role of nuclear receptor Nur77 in *Escherichia coli*-induced peritonitis. [Infect Immun. 82 \(1\): 253-64.](#)
15. Roche, J.A. *et al.* (2015) Myofiber damage precedes macrophage infiltration after *in vivo* injury in dysferlin-deficient a/j mouse skeletal muscle. [Am J Pathol. 185 \(6\): 1686-98.](#)
16. Lee, Y.S. *et al.* (2015) Interleukin-1 (IL-1) signaling in intestinal stromal cells controls KC/ CXCL1 secretion, which correlates with recruitment of IL-22- secreting neutrophils at early stages of *Citrobacter rodentium* infection. [Infect Immun. 83 \(8\): 3257-67.](#)
17. Heckelsmiller, K. *et al.* (2002) Combined dendritic cell- and CpG oligonucleotide-based immune therapy cures large murine tumors that resist chemotherapy. [Eur J Immunol. 32 \(11\): 3235-45.](#)
18. Zhang, M.Z. *et al.* (2015) Inhibition of cyclooxygenase-2 in hematopoietic cells results in salt-sensitive hypertension. [J Clin Invest. 125 \(11\): 4281-94.](#)
19. Leblond, A.L. *et al.* (2015) Systemic and Cardiac Depletion of M2 Macrophage through CSF-1R Signaling Inhibition Alters Cardiac Function Post Myocardial Infarction. [PLoS One. 10 \(9\): e0137515.](#)
20. Kojo, K. *et al.* (2016) BLT1 signalling protects the liver against acetaminophen hepatotoxicity by preventing excessive accumulation of hepatic neutrophils. [Sci Rep. 6: 29650.](#)
21. Otsuru, S. *et al.* (2017) Hematopoietic derived cells do not contribute to osteogenesis as osteoblasts. [Bone. 94: 1-9.](#)
22. Wang, Y. *et al.* (2015) Proximal tubule-derived colony stimulating factor-1 mediates polarization of renal macrophages and dendritic cells, and recovery in acute kidney injury. [Kidney Int. 88 \(6\): 1274-1282.](#)
23. Cousins, F.L. *et al.* (2016) Evidence for a dynamic role for mononuclear phagocytes during endometrial repair and remodelling. [Sci Rep. 6: 36748.](#)
24. Cotrina ML *et al.* (2017) Direct comparison of microglial dynamics and inflammatory profile in photothrombotic and arterial occlusion evoked stroke. [Neuroscience. 343: 483-94.](#)
25. Kamata, M. *et al.* (2019) Role of the high-affinity leukotriene B₄ receptor signaling in

- fibrosis after unilateral ureteral obstruction in mice. [PLoS One. 14 \(2\): e0202842.](#)
26. Natanov, R. *et al.* (2018) Blood cytokine expression correlates with early multi-organ damage in a mouse model of moderate hypothermia with circulatory arrest using cardiopulmonary bypass. [PLoS One. 13 \(10\): e0205437.](#)
27. Idowu, T.O. *et al.* (2020) Identification of specific Tie2 cleavage sites and therapeutic modulation in experimental sepsis. [Elife. 9: e59520.](#)
28. Qin, X. *et al.* (2018) Caspase recruitment domain-containing protein 9 (CARD9) knockout reduces regional ischemia/reperfusion injury through an attenuated inflammatory response. [PLoS One. 13 \(6\): e0199711.](#)
29. Takahashi, R. *et al.* (2020) Microsomal prostaglandin E synthase-1 promotes lung metastasis via SDF-1/CXCR4-mediated recruitment of CD11b⁺Gr1⁺MDSCs from bone marrow. [Biomed Pharmacother. 121: 109581.](#)
30. Idowu, T.O. *et al.* (2021) Flow-dependent regulation of endothelial Tie2 by GATA3 *in vivo.* [Intensive Care Med Exp. 9 \(1\): 38.](#)
31. Zhang, M.Z. *et al.* (2019) The Role of the EGF Receptor in Sex Differences in Kidney Injury. [J Am Soc Nephrol. 30 \(9\): 1659-73.](#)
32. Schünke, H. *et al.* (2021) OTULIN inhibits RIPK1-mediated keratinocyte necroptosis to prevent skin inflammation in mice. [Nat Commun. 12 \(1\): 5912.](#)
33. Konishi, T. *et al.* (2019) Cell-specific regulatory effects of CXCR2 on cholestatic liver injury. [Am J Physiol Gastrointest Liver Physiol. 317 \(6\): G773-G783.](#)
34. Christodoulou-Vafeiadou, E. *et al.* (2020) Ectopic bone formation and systemic bone loss in a transmembrane TNF-driven model of human spondyloarthritis. [Arthritis Res Ther. 22 \(1\): 232.](#)
35. Störmer, J. *et al.* (2022) A Single Oral Dose of Diclofenac Causes Transition of Experimental Subclinical Acute Kidney Injury to Chronic Kidney Disease. [Biomedicines. 10 \(5\): 1198.](#)
36. Greite, R. *et al.* (2022) Cell-Free Hemoglobin in Acute Kidney Injury after Lung Transplantation and Experimental Renal Ischemia/Reperfusion. [Int J Mol Sci. 23\(21\):13272.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®550 , DyLight®650 , DyLight®800
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M414840:221214'

Printed on 23 Feb 2023