

Datasheet: MCA2388GA

BATCH NUMBER 167887

Description:	RAT ANTI MOUSE CD31
Specificity:	CD31
Other names:	PECAM-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	ER-MP12
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/100
Immunohistology - Frozen (1)	▪			
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	BALB/c macrophage precursor cell hybrids
External Database Links	<p>UniProt: Q08481 Related reagents</p> <p>Entrez Gene: 18613 Pecam1 Related reagents</p>
Synonyms	Pecam, Pecam-1
RRID	AB_2161024
Fusion Partners	Cells from immunised rats were fused with the cells of the rat Y3-Ag1.2.3 myeloma cell line
Specificity	<p>Rat anti Mouse CD31 antibody, clone ER-MP12 recognizes mouse CD31, a 140 kDa cell surface glycoprotein expressed at high levels on endothelial cells, platelets and most leukocyte subpopulations.</p> <p>CD31 is also expressed on a major population of macrophage / dendritic cell precursors in the bone marrow. Rat anti Mouse CD31 antibody, clone ER-MP12 can be used in conjunction with clone ER-MP20 (MCA2389GA) in two colour flow cytometric analysis, to identify different stages of myeloid progenitor cells in mouse bone marrow (de Bruijn <i>et al.</i> 1998).</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> 1. Leenen, P.J. <i>et al.</i> (1990) Murine macrophage precursor characterization. II. Monoclonal antibodies against macrophage precursor antigens. Eur J Immunol. 20 (1): 27-34. 2. van der Loo, J. <i>et al.</i> (1995) Identification of hematopoietic stem cell subsets on the basis of their primitiveness using antibody ER-MP12. Blood. 85:952-62. 3. Ling, V. <i>et al.</i> (1997) Structural identification of the hematopoietic progenitor antigen ER-MP12 as the vascular endothelial adhesion molecule PECAM-1 (CD31). Eur J Immunol. 27:509-14. 4. de Bruijn, M.F. <i>et al.</i> (1998) Bone marrow cellular composition in Listeria monocytogenes infected mice detected using ER-MP12 and ER-MP20 antibodies: a flow cytometric alternative to differential counting. J Immunol Methods. 217 (1-2): 27-39. 5. Wynn, A.A. <i>et al.</i> (2001) Role of granulocyte/macrophage colony-stimulating factor in

- zymocel-induced hepatic granuloma formation. [Am J Pathol. 158 \(1\): 131-45.](#)
6. van Rijt, L. *et al.* (2002) Allergen-induced accumulation of airway dendritic cells is supported by an increase in CD31(hi)Ly-6C(neg) bone marrow precursors in a mouse model of asthma. [Blood. 100:3663-71.](#)
 7. Tagoh, H. *et al.* (2002) Transcription factor complex formation and chromatin fine structure alterations at the murine c-fms (CSF-1 receptor) locus during maturation of myeloid precursor cells. [Genes Dev. 16:1721-37.](#)
 8. Nikolic, T. *et al.* (2002) Developmental stages of myeloid dendritic cells in mouse bone marrow. [Int Immunol. 15:515-24.](#)
 9. Baumeister, T. *et al.* (2003) Interleukin-3Ralpha+ myeloid dendritic cells and mast cells develop simultaneously from different bone marrow precursors in cultures with interleukin-3. [J Invest Dermatol. 121: 280-8.](#)
 10. Geutskens, S.B. *et al.* (2005) Macrophages in the murine pancreas and their involvement in fetal endocrine development *in vitro*. [J Leukoc Biol. 78: 845-52.](#)
 11. Revermann, M. *et al.* (2010) Soluble epoxide hydrolase deficiency attenuates neointima formation in the femoral cuff model of hyperlipidemic mice. [Arterioscler Thromb Vasc Biol. 30: 909-14.](#)
 12. Sumagin, R. and Sarelius, I.H. (2010) Intercellular adhesion molecule-1 enrichment near tricellular endothelial junctions is preferentially associated with leukocyte transmigration and signals for reorganization of these junctions to accommodate leukocyte passage. [J Immunol. 184: 5242-52.](#)
 13. Ross, E.A. *et al.* (2011) CD31 Is Required on CD4+ T Cells To Promote T Cell Survival during *Salmonella* Infection. [J Immunol. 187: 1553-65.](#)
 14. Thum, T. *et al.* (2011) Impairment of endothelial progenitor cell function and vascularization capacity by aldosterone in mice and humans. [Eur Heart J. 32: 1275-86.](#)
 15. Thorp, E. *et al.* (2011) A reporter for tracking the UPR *in vivo* reveals patterns of temporal and cellular stress during atherosclerotic progression. [J Lipid Res. 52 \(5\): 1033-8.](#)
 16. Schledzewski, K. *et al.* (2011) Deficiency of liver sinusoidal scavenger receptors stabilin-1 and -2 in mice causes glomerulofibrotic nephropathy via impaired hepatic clearance of noxious blood factors. [J Clin Invest. 121: 703-14.](#)
 17. Loureiro, J. *et al.* (2011) Blocking TGF- β 1 Protects the Peritoneal Membrane from Dialysate-Induced Damage. [J Am Soc Nephrol. 22: 1682-95.](#)
 18. Matsakas, A. *et al.* (2012) Exercise training attenuates the hypermuscular phenotype and restores skeletal muscle function in the myostatin null mouse. [Exp Physiol. 97 \(1\): 125-40.](#)
 19. Trottier MD *et al.* (2012) Enhancement of hematopoiesis and lymphopoiesis in diet-induced obese mice. [Proc Natl Acad Sci U S A. 109 \(20\): 7622-9.](#)
 20. Moen, I. *et al.* (2012) Gene expression in tumor cells and stroma in dsRed 4T1 tumors in eGFP-expressing mice with and without enhanced oxygenation. [BMC Cancer. 12: 21.](#)
 21. Trottier, M.D. *et al.* (2012) Enhanced production of early lineages of monocytic and granulocytic cells in mice with colitis [Proc Natl Acad Sci U S A. 109: 16594-9.](#)
 22. Kroon, P. *et al.* (2013) JAK-STAT blockade inhibits tumor initiation and clonogenic recovery of prostate cancer stem-like cells. [Cancer Res. 73 \(16\): 5288-98.](#)
 23. Ono, N. *et al.* (2014) A subset of chondrogenic cells provides early mesenchymal progenitors in growing bones. [Nat Cell Biol. 16 \(12\): 1157-67.](#)
 24. Eskilsson, A. *et al.* (2014) Distribution of microsomal prostaglandin E synthase-1 in the

- mouse brain. [J Comp Neurol. 522 \(14\): 3229-44.](#)
25. Nakamura, Y. *et al.* (2015) Mesenchymal-stem-cell-derived exosomes accelerate skeletal muscle regeneration. [FEBS Lett. 589 \(11\): 1257-65.](#)
26. Fraccarollo, D. *et al.* (2015) Efficacy of mineralocorticoid receptor antagonism in the acute myocardial infarction phase: eplerenone versus spironolactone. [ESC Heart Fail. 2 \(3\): 150-8.](#)
27. Stein-Merlob, A.F. *et al.* (2015) Blood Accessibility to Fibrin in Venous Thrombosis is Thrombus Age-Dependent and Predicts Fibrinolytic Efficacy: An *In Vivo* Fibrin Molecular Imaging Study. [Theranostics. 5 \(12\): 1317-27.](#)
28. Cao Y *et al.* (2016) IL-1 β differently stimulates proliferation and multinucleation of distinct mouse bone marrow osteoclast precursor subsets. [J Leukoc Biol. 100 \(3\): 513-23.](#)
29. Ryan, T.E. *et al.* (2016) Mitochondrial therapy improves limb perfusion and myopathy following hindlimb ischemia. [J Mol Cell Cardiol. 97: 191-6.](#)
30. Shi, H. *et al.* (2016) Hiding inside? Intracellular expression of non-glycosylated c-kit protein in cardiac progenitor cells. [Stem Cell Res. 16 \(3\): 795-806.](#)
31. Chowdhury, B. *et al.* (2016) Hyaluronidase 2 (HYAL2) is expressed in endothelial cells, as well as some specialized epithelial cells, and is required for normal hyaluronan catabolism. [Histochem Cell Biol. 145 \(1\): 53-66.](#)
32. Yip, H.K. *et al.* (2016) Tissue plasminogen activator deficiency preserves neurological function and protects against murine acute ischemic stroke. [Int J Cardiol. 205: 133-41.](#)
33. Reigstad, I. *et al.* (2016) The Effect of Stromal Integrin β 3-Deficiency on Two Different Tumors in Mice. [Cancers \(Basel\). 8 \(1\): pii: E14.](#)
34. Bongiorno, E.K. *et al.* (2017) Type 1 Immune Mechanisms Driven by the Response to Infection with Attenuated Rabies Virus Result in Changes in the Immune Bias of the Tumor Microenvironment and Necrosis of Mouse GL261 Brain Tumors. [J Immunol. 198 \(11\): 4513-4523.](#)
35. Cao, Y. *et al.* (2017) TNF- α has both stimulatory and inhibitory effects on mouse monocyte-derived osteoclastogenesis. [J Cell Physiol. 232 \(12\): 3273-85.](#)
36. Piro, J.R. *et al.* (2018) Inhibition of 2-AG hydrolysis differentially regulates blood brain barrier permeability after injury. [J Neuroinflammation. 15 \(1\): 142.](#)
37. Oikawa, S. *et al.* (2018) Role of endothelial microRNA-23 clusters in angiogenesis in vivo. [Am J Physiol Heart Circ Physiol. 315 \(4\): H838-H846.](#)
38. Tay, M.H.D. *et al.* (2019) Halted Lymphocyte Egress via Efferent Lymph Contributes to Lymph Node Hypertrophy During Hypercholesterolemia. [Front Immunol. 10: 575.](#)
39. Iring, A. *et al.* (2019) Shear stress-induced endothelial adrenomedullin signaling regulates vascular tone and blood pressure. [J Clin Invest. 129 \(7\): 2775-91.](#)
40. Ascone, G. *et al.* (2020) Increase in the Number of Bone Marrow Osteoclast Precursors at Different Skeletal Sites, Particularly in Long Bone and Jaw Marrow in Mice Lacking IL-1RA. [Int J Mol Sci. 21\(11\):3774.](#)
41. Patel, P. *et al.* (2023) Development of a Personalised Device for Systemic Magnetic Drug Targeting to Brain Tumours. [Nanotheranostics. 7 \(1\): 102-16.](#)
42. Eskilsson, A. *et al.* (2023) Prostaglandin production in brain endothelial cells during the initiation of fever. [Commun Integr Biol. 16 \(1\): 2166237.](#)
43. Yamashita, J. *et al.* (2021) Effect of bisphosphonates on healing of tooth extraction wounds in infectious osteomyelitis of the jaw. [Bone. 143: 115611.](#)
44. Molás, R.B. *et al.* (2020) The involvement of annexin A1 in human placental response to maternal Zika virus infection. [Antiviral Res. 179: 104809.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2388GA 10040
--------------------------------------	---

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®550 , DyLight®650 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M414856:221214'

Printed on 29 Feb 2024