

Datasheet: MCA2389GA

BATCH NUMBER 169292

Description:	RAT ANTI MOUSE Ly-6C
Specificity:	Ly-6C
Other names:	Lymphocyte antigen 6C2
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	ER-MP20
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/50
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃)

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Balb/c macrophage precursor cell hybrids.
External Database Links	UniProt: P0CW03 Related reagents
RRID	AB_844551
Fusion Partners	Spleen cells from immunized rats were fused with cells of the Y3-Ag1.2.3 myeloma cell line.
Specificity	<p>Rat anti Mouse Ly-6C antibody, clone ER-MP20 recognizes murine Ly-6C, a 131 amino acid ~14 kDa differentiation antigen, expressed on macrophage/dendritic cell precursors in mid-stage development (late CFU-M, monoblasts and immature monocytes), granulocytes, and on a wide range of endothelial cells and subpopulations of B- and T-lymphocytes.</p> <p>Rat anti Mouse Ly-6C antibody, clone ER-MP20 is able to distinguish multiple mouse blood monocyte subsets: immature Ly-6C^{hi} monocytes are recruited to acute peripheral inflammation and develop into Ly-6C⁺ exudate macrophages, whereas more mature Ly-6C^{lo} monocytes are precursors for tissue macrophages and dendritic cells in steady state.</p> <p>Rat anti Mouse Ly-6C, clone ER-MP20 can be used in conjunction with clone ER-MP12 in two colour flow cytometric analysis, to identify different stages of myeloid progenitor cells in mouse bone marrow (Leenen et al. 1990).</p> <p>Rat anti Mouse Ly-6C was originally described as recognizing a protein encoded by the LY6C gene. It has subsequently become apparent that the LY6C locus demonstrates polymorphism and the LY6C gene has been re-designated LY6C2. The LY6C1 gene encodes a similar protein with ~95% sequence homology to LY6C2.</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> Zhang, Y. & Bliska, J.B. (2010) YopJ-promoted cytotoxicity and systemic colonization are associated with high levels of murine interleukin-18, gamma interferon, and neutrophils in a live vaccine model of <i>Yersinia pseudotuberculosis</i> infection. Infect Immun 78: 2329-41. Leenen, P.J. et al. (1990) Murine macrophage precursor characterization. II. Monoclonal antibodies against macrophage precursor antigens. Eur J Immunol. 20 (1): 27-34. de Bruijn, M.F. et al. (1998) Bone marrow cellular composition in Listeria monocytogenes infected mice detected using ER-MP12 and ER-MP20 antibodies: a flow cytometric alternative to differential counting. J Immunol Methods. 217 (1-2): 27-39.

4. Schatteman, G.C. *et al.* (2010) Lin- Cells Mediate Tissue Repair by Regulating MCP-1/CCL-2. [Am J Pathol. 177: 2002-10.](#)
5. Baumeister, T. *et al.* (2003) Interleukin-3 α myeloid dendritic cells and mast cells develop simultaneously from different bone marrow precursors in cultures with interleukin-3. [J Invest Dermatol. 121: 280-8.](#)
6. Devey, L. *et al.* (2009) Tissue-resident macrophages protect the liver from ischemia reperfusion injury via a heme oxygenase-1-dependent mechanism. [Mol Ther. 17: 65-72.](#)
7. Nikolic, T. *et al.* (2003) Developmental stages of myeloid dendritic cells in mouse bone marrow. [Int Immunol. 15: 515-24.](#)
8. Wynn, A.A. *et al.* (2001) Role of granulocyte/macrophage colony-stimulating factor in zymocel-induced hepatic granuloma formation. [Am J Pathol. 158 \(1\): 131-45.](#)
9. Lesokhin, A.M. *et al.* (2012) Monocytic CCR2+ Myeloid-Derived Suppressor Cells Promote Immune Escape by Limiting Activated CD8 T-cell Infiltration into the Tumor Microenvironment. [Cancer Res. 72: 876-86.](#)
10. Chan, J. *et al.* (1998) Macrophage lineage cells in inflammation: characterization by colony-stimulating factor-1 (CSF-1) receptor (c-Fms), ER-MP58, and ER-MP20 (Ly-6C) expression. [Blood. 92: 1423-31.](#)
11. van Rijt, L.S. *et al.* (2002) Allergen-induced accumulation of airway dendritic cells is supported by an increase in CD31(hi)Ly-6C(neg) bone marrow precursors in a mouse model of asthma. [Blood. 100: 3663-71.](#)
12. Arnardottir, H.H.*et al.* (2012) Dietary Fish Oil Decreases the Proportion of Classical Monocytes in Blood in Healthy Mice but Increases Their Proportion upon Induction of Inflammation. [J Nutr. 142: 803-8.](#)
13. Henkel, G. *et al.* (1999) Commitment to the monocytic lineage occurs in the absence of the transcription factor PU.1. [Blood. 93:2849-58.](#)
14. Bossaller, L. *et al.* (2013) Overexpression of membrane-bound fas ligand (CD95L) exacerbates autoimmune disease and renal pathology in pristane-induced lupus. [J Immunol. 191: 2104-14.](#)
15. Garcia, J.A. *et al.* (2013) Regulation of adaptive immunity by the fractalkine receptor during autoimmune inflammation. [J Immunol. 191: 1063-72.](#)
16. Stijlemans, B. *et al.* (2015) Murine Liver Myeloid Cell Isolation Protocol [BIO-PROTOCOL. 5 \(10\).](#)
17. Damya, L. *et al.* (2014) Purification of Tumor-Associated Macrophages (TAM) and Tumor-Associated Dendritic Cells (TADC) [BIO-PROTOCOL. 4 \(22\).](#)
18. Morganti, J.M. *et al.* (2016) Age exacerbates the CCR2/5-mediated neuroinflammatory response to traumatic brain injury. [J Neuroinflammation. 13 \(1\): 80.](#)
19. Mooney, J.E. *et al.* (2010) Cellular plasticity of inflammatory myeloid cells in the peritoneal foreign body response. [Am J Pathol. 176 \(1\): 369-80.](#)
20. Iwasaki, Y. *et al.* (2011) *In situ* proliferation and differentiation of macrophages in dental pulp. [Cell Tissue Res. 346 \(1\): 99-109.](#)
21. Movahedi, K. *et al.* (2012) Nanobody-based targeting of the macrophage mannose receptor for effective in vivo imaging of tumor-associated macrophages. [Cancer Res. 72 \(16\): 4165-77.](#)
22. Ribechini, E. *et al.* (2009) Gr-1 antibody induces STAT signaling, macrophage marker expression and abrogation of myeloid-derived suppressor cell activity in BM cells. [Eur J Immunol. 39 \(12\): 3538-51.](#)
23. Bossaller, L. *et al.* (2016) TLR9 Deficiency Leads to Accelerated Renal Disease and

- Myeloid Lineage Abnormalities in Pristane-Induced Murine Lupus. [J Immunol. 197 \(4\): 1044-53.](#)
24. Barnes, M.A. *et al.* (2015) Macrophage migration inhibitory factor is required for recruitment of scar-associated macrophages during liver fibrosis. [J Leukoc Biol. 97 \(1\): 161-9.](#)
25. Ohnishi, K. *et al.* (2012) Immunohistochemical detection of possible cellular origin of hepatic histiocytic sarcoma in mice. [J Clin Exp Hematop. 52 \(3\): 171-7.](#)
26. Van den Bossche. J. *et al.* (2012) Claudin-1, claudin-2 and claudin-11 genes differentially associate with distinct types of anti-inflammatory macrophages *in vitro* and with parasite- and tumour-elicited macrophages *in vivo*. [Scand J Immunol. 75 \(6\): 588-98.](#)
27. Houthuys, E. *et al.* (2010) A method for the isolation and purification of mouse peripheral blood monocytes. [J Immunol Methods. 359 \(1-2\): 1-10.](#)
28. Greifenberg, V. *et al.* (2009) Myeloid-derived suppressor cell activation by combined LPS and IFN-gamma treatment impairs DC development. [Eur J Immunol. 39 \(10\): 2865-76.](#)
29. Cardona, S.M. *et al.* (2015) Disruption of Fractalkine Signaling Leads to Microglial Activation and Neuronal Damage in the Diabetic Retina. [ASN Neuro. 7 \(5\):1759091415608204.](#)
30. Waddell, A. *et al.* (2011) Colonic eosinophilic inflammation in experimental colitis is mediated by Ly6C(high) CCR2(+) inflammatory monocyte/macrophage-derived CCL11. [J Immunol. 186 \(10\): 5993-6003.](#)
31. Robbie, S.J. *et al.* (2016) Enhanced Ccl2-Ccr2 signaling drives more severe choroidal neovascularization with aging. [Neurobiol Aging. 40: 110-9.](#)
32. Cao, Y. *et al.* (2016) IL-1 β differently stimulates proliferation and multinucleation of distinct mouse bone marrow osteoclast precursor subsets. [J Leukoc Biol. 100 \(3\): 513-23.](#)
33. Cao, Y. *et al.* (2017) TNF- α has both stimulatory and inhibitory effects on mouse monocyte-derived osteoclastogenesis. [J Cell Physiol. 232 \(12\): 3273-85.](#)
34. Khedoe, P.P.S.J. *et al.* (2017) Acute and chronic effects of treatment with mesenchymal stromal cells on LPS-induced pulmonary inflammation, emphysema and atherosclerosis development. [PLoS One. 12 \(9\): e0183741.](#)
35. Koohy, H. *et al.* (2018) Genome organization and chromatin analysis identify transcriptional downregulation of insulin-like growth factor signaling as a hallmark of aging in developing B cells. [Genome Biol. 19 \(1\): 126.](#)
36. Pluijmer, N.J. *et al.* (2020) Effects on cardiac function, remodeling and inflammation following myocardial ischemia-reperfusion injury or unperfused myocardial infarction in hypercholesterolemic APOE*3-Leiden mice. [Sci Rep. 10 \(1\): 16601.](#)
37. Ascone, G. *et al.* (2020) Increase in the Number of Bone Marrow Osteoclast Precursors at Different Skeletal Sites, Particularly in Long Bone and Jaw Marrow in Mice Lacking IL-1RA. [Int J Mol Sci. 21 \(11\): 3774.](#)
38. Pluijmer, N.J. *et al.* (2021) Phosphorylcholine antibodies restrict infarct size and left ventricular remodelling by attenuating the unperfused post-ischaemic inflammatory response. [J Cell Mol Med. 25 \(16\): 7772-82.](#)
39. Njock, M-K. (2022) Endothelial extracellular vesicles promote tumour growth by tumour-associated macrophage reprogramming [J Extracell Vesicles 2022 Jun;11\(6\):e12228.](#)
40. Vainchtein, I.D. *et al.* (2023) Characterizing microglial gene expression in a model of secondary progressive multiple sclerosis. [Glia. 71 \(3\): 588-601.](#)

41. Mielczarek, O. *et al.* (2023) Intra- and interchromosomal contact mapping reveals the Igh locus has extensive conformational heterogeneity and interacts with B-lineage genes. [Cell Rep 42 \(9\):113074.](#)
42. Thiem, K. *et al.* (2019) Deletion of hematopoietic Dectin-2 or CARD9 does not protect against atherosclerotic plaque formation in hyperlipidemic mice. [Sci Rep. 9 \(1\): 4337.](#)
43. Heissig, B. *et al.* (2022) siRNA against CD40 delivered via a fungal recognition receptor ameliorates murine acute graft-versus-host disease. [EJHaem. 3 \(3\): 849-61.](#)
44. Cardona, S.M. *et al.* (2018) Role of the Fractalkine Receptor in CNS Autoimmune Inflammation: New Approach Utilizing a Mouse Model Expressing the Human CX3CR1(I249/M280) Variant. [Front Cell Neurosci. 12: 365.](#)
45. Burzynski, L.C. *et al.* (2023) Thrombin-activated interleukin-1 α drives atherogenesis, but also promotes vascular smooth muscle cell proliferation and collagen production. [Cardiovasc Res. 119 \(12\): 2179-89.](#)
46. Story, M.E. *et al.* (2024) Resident memory T cells in dirty mice suppress innate cell activation and infiltration into the skin following stimulation with alarmins [BioRxiv 16 Jul \[Epub ahead of print\].](#)
47. Lepland, A. *et al.* (2024) Therapeutic Tumor Macrophage Reprogramming in Breast Cancer Through a Peptide-Drug Conjugate [bioRxiv: 12 Aug. \[Epub ahead of print\].](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2389GA 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight@800
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight@550 , DyLight@650 , DyLight@800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR72...)	HRP

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M414893:221214'

Printed on 12 Dec 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)