

Datasheet: MCA2389SBUV445 BATCH NUMBER 100006496

Description:	RAT ANTI MOUSE Ly-6C:StarBright UltraViolet 445		
Specificity:	Ly-6C		
Other names:	Lymphocyte antigen 6C2		
Format:	StarBright UltraViolet 445		
Product Type:	Monoclonal Antibody		
Clone:	ER-MP20		
Isotype:	lgG2a		
Quantity:	100 TESTS/0.5ml		

Product Details

Applications	This product has been reported to work in the following applications. This info derived from testing within our laboratories, peer-reviewed publications or per communications from the originators. Please refer to references indicated for information. For general protocol recommendations, please visit <u>www.bio-</u> rad-antibodies.com/protocols.				
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry	-			Neat
	Where this product ha	as not been test	ed for u	ise in a particular tech	nnique this does not
	necessarily exclude it a guide only. It is reco system using appropr	mmended that	the use	r titrates the product	ng dilutions are given as for use in their own
Target Species	Mouse				
Product Form	Purified IgG conjugated to StarBright UltraViolet 445 - liquid				
Max Ex/Em	Fluorophore	Excitation Max	(nm)	Emission Max (nm)	
	StarBright UltraViolet 445	347		440	
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant				
Buffer Solution	Phosphate buffered saline				
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Alb 0.1% Pluronic F68				

0.1% PEG 3350 0.05% Tween 20

Balb/c macrophage precursor cell hybrids.
UniProt: <u>P0CW03</u> <u>Related reagents</u>
Spleen cells from immunised rats were fused with cells of the Y3-Ag1.2.3 myeloma cell line.
Rat anti Mouse Ly-6C antibody, clone ER-MP20 recognizes murine Ly-6C, a 131 amino acid ~14 kDa differentiation antigen, expressed on macrophage/dendritic cell precursors in mid-stage development (late CFU-M, monoblasts and immature monocytes), granulocytes, and on a wide range of endothelial cells and subpopulations of B- and T-lymphocytes.
Rat anti Mouse Ly-6C antibody, clone ER-MP20 is able to distinguish multiple mouse blood monocyte subsets: immature Ly-6C ^{hi} monocytes are recruited to acute peripheral inflammation and develop into Ly-6C ⁺ exudate macrophages, whereas more mature Ly-6C ^{-/lo} monocytes are precursors for tissue macrophages and dendritic cells in steady state.
Rat anti Mouse Ly-6C, clone ER-MP20 can be used in conjunction with clone <u>ER-MP12</u> in two colour flow cytometric analysis, to identify different stages of myeloid progenitor cells in mouse bone marrow (<u>Leenen <i>et al.</i> 1990</u>).
Rat anti Mouse Ly-6C was originally described as recognizing a protein encoded by the LY6C gene. It has subsequently become apparent that the LY6C locus demonstrates polymorphism and the LY6C gene has been re-designated LY6C2. The LY6C1 gene encodes a similar protein with ~95% sequence homology to LY6C2.
Use 5ul of the suggested working dilution to label 10 ⁶ cells in 100ul. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
 Zhang, Y. & Bliska, J.B. (2010) YopJ-promoted cytotoxicity and systemic colonization are associated with high levels of murine interleukin-18, gamma interferon, and neutrophils in a live vaccine model of <i>Yersinia pseudotuberculosis</i> infection. Infect Immun 78: 2329-41. Leenen, P.J. <i>et al.</i> (1990) Murine macrophage precursor characterization. II. Monoclonal antibodies against macrophage precursor antigens. <u>Eur J Immunol. 20 (1): 27-34.</u> de Bruijn, M.F. <i>et al.</i> (1998) Bone marrow cellular composition in Listeria monocytogenes infected mice detected using ER-MP12 and ER-MP20 antibodies: a flow cytometric alternative to differential counting. <u>J Immunol Methods. 217 (1-2): 27-39.</u> Schatteman, G.C. <i>et al.</i> (2010) Lin- Cells Mediate Tissue Repair by Regulating MCP-1/CCL-2. <u>Am J Pathol. 177: 2002-10.</u> Baumeister, T. <i>et al.</i> (2003) Interleukin-3Ralpha+ myeloid dendritic cells and mast cells

develop simultaneously from different bone marrow precursors in cultures with interleukin-3. J Invest Dermatol. 121: 280-8.

Devey, L. *et al.* (2009) Tissue-resident macrophages protect the liver from ischemia reperfusion injury via a heme oxygenase-1-dependent mechanism. <u>Mol Ther. 17: 65-72.</u>
 Nikolic, T. *et al.* (2003) Developmental stages of myeloid dendritic cells in mouse bone marrow. <u>Int Immunol. 15: 515-24.</u>

8. Wynn, A.A. *et al.* (2001) Role of granulocyte/macrophage colony-stimulating factor in zymocel-induced hepatic granuloma formation. <u>Am J Pathol. 158 (1): 131-45.</u>

9. Lesokhin, A.M. *et al.* (2012) Monocytic CCR2+ Myeloid-Derived Suppressor Cells Promote Immune Escape by Limiting Activated CD8 T-cell Infiltration into the Tumor Microenvironment. <u>Cancer Res. 72: 876-86.</u>

10. Chan, J. *et al.* (1998) Macrophage lineage cells in inflammation: characterization by colony-stimulating factor-1 (CSF-1) receptor (c-Fms), ER-MP58, and ER-MP20 (Ly-6C) expression. <u>Blood. 92: 1423-31.</u>

11. van Rijt, L.S. *et al.* (2002) Allergen-induced accumulation of airway dendritic cells is supported by an increase in CD31(hi)Ly-6C(neg) bone marrow precursors in a mouse model of asthma. <u>Blood. 100: 3663-71.</u>

12. Arnardottir, H.H.*et al.* (2012) Dietary Fish Oil Decreases the Proportion of Classical Monocytes in Blood in Healthy Mice but Increases Their Proportion upon Induction of Inflammation. <u>J Nutr. 142: 803-8.</u>

13. Henkel, G. *et al.* (1999) Commitment to the monocytic lineage occurs in the absence of the transcription factor PU.1. <u>Blood. 93:2849-58.</u>

14. Bossaller, L. *et al.* (2013) Overexpression of membrane-bound fas ligand (CD95L) exacerbates autoimmune disease and renal pathology in pristane-induced lupus. J Immunol. 191: 2104-14.

15. Garcia, J.A. *et al.* (2013) Regulation of adaptive immunity by the fractalkine receptor during autoimmune inflammation. <u>J Immunol. 191: 1063-72.</u>

16. Benoit, S. *et al.* (2015) Murine Liver Myeloid Cell Isolation Protocol <u>BIO-PROTOCOL.</u> <u>5 (10) [Epub ahead of print].</u>

17. Damya, L. *et al.* (2014) Purification of Tumor-Associated Macrophages (TAM) and Tumor-Associated Dendritic Cells (TADC) <u>BIO-PROTOCOL. 4 (22) [Epub ahead of print].</u>

18. Morganti, J.M. *et al.* (2016) Age exacerbates the CCR2/5-mediated neuroinflammatory response to traumatic brain injury. <u>J Neuroinflammation. 13 (1): 80.</u>

19. Mooney, J.E. *et al.* (2010) Cellular plasticity of inflammatory myeloid cells in the peritoneal foreign body response. <u>Am J Pathol. 176 (1): 369-80.</u>

20. Iwasaki, Y. *et al.* (2011) *In situ* proliferation and differentiation of macrophages in dental pulp. <u>Cell Tissue Res. 346 (1): 99-109.</u>

21. Movahedi, K. *et al.* (2012) Nanobody-based targeting of the macrophage mannose receptor for effective in vivo imaging of tumor-associated macrophages. <u>Cancer Res. 72</u> (<u>16</u>): <u>4165-77</u>.

22. Ribechini, E. *et al.* (2009) Gr-1 antibody induces STAT signaling, macrophage marker expression and abrogation of myeloid-derived suppressor cell activity in BM cells. <u>Eur J</u> <u>Immunol. 39 (12): 3538-51.</u>

23. Bossaller, L. *et al.* (2016) TLR9 Deficiency Leads to Accelerated Renal Disease and Myeloid Lineage Abnormalities in Pristane-Induced Murine Lupus. <u>J Immunol. 197 (4):</u> 1044-53.

24. Barnes, M.A. et al. (2015) Macrophage migration inhibitory factor is required for

recruitment of scar-associated macrophages during liver fibrosis. <u>J Leukoc Biol. 97 (1):</u> <u>161-9.</u>

25. Ohnishi, K. *et al.* (2012) Immunohistochemical detection of possible cellular origin of hepatic histiocytic sarcoma in mice. J Clin Exp Hematop. 52 (3): 171-7.

26. Van den Bossche. J. *et al.* (2012) Claudin-1, claudin-2 and claudin-11 genes differentially associate with distinct types of anti-inflammatory macrophages *in vitro* and with parasite- and tumour-elicited macrophages *in vivo*. <u>Scand J Immunol. 75 (6): 588-98.</u>

27. Houthuys, E. *et al.* (2010) A method for the isolation and purification of mouse peripheral blood monocytes. J Immunol Methods. 359 (1-2): 1-10.

28. Greifenberg, V. *et al.* (2009) Myeloid-derived suppressor cell activation by combined LPS and IFN-gamma treatment impairs DC development. <u>Eur J Immunol. 39 (10):</u> 2865-76.

29. Cardona, S.M.*et al.* (2015) Disruption of Fractalkine Signaling Leads to Microglial Activation and Neuronal Damage in the Diabetic Retina. <u>ASN Neuro. 7 (5)Oct 29 [Epub ahead of print].</u>

30. Waddell, A. *et al.* (2011) Colonic eosinophilic inflammation in experimental colitis is mediated by Ly6C(high) CCR2(+) inflammatory monocyte/macrophage-derived CCL11. J Immunol. 186 (10): 5993-6003.

31. Robbie, S.J. *et al.* (2016) Enhanced Ccl2-Ccr2 signaling drives more severe choroidal neovascularization with aging. <u>Neurobiol Aging. 40: 110-9.</u>

32. Cao, Y. *et al.* (2016) IL-1 β differently stimulates proliferation and multinucleation of distinct mouse bone marrow osteoclast precursor subsets. <u>J Leukoc Biol. 100 (3): 513-23</u>. 33. Cao, Y. *et al.* (2017) TNF- α has both stimulatory and inhibitory effects on mouse monocyte-derived osteoclastogenesis. <u>J Cell Physiol. 232 (12): 3273-85</u>.

34. Khedoe, P.P.S.J. *et al.* (2017) Acute and chronic effects of treatment with mesenchymal stromal cells on LPS-induced pulmonary inflammation, emphysema and atherosclerosis development. <u>PLoS One. 12 (9): e0183741</u>.

35. Koohy, H. *et al.* (2018) Genome organization and chromatin analysis identify transcriptional downregulation of insulin-like growth factor signaling as a hallmark of aging in developing B cells. <u>Genome Biol. 19 (1): 126.</u>

36. Pluijmert, N.J. *et al.* (2020) Effects on cardiac function, remodeling and inflammation following myocardial ischemia-reperfusion injury or unreperfused myocardial infarction in hypercholesterolemic APOE*3-Leiden mice. <u>Sci Rep. 10 (1): 16601.</u>

37. Ascone, G. *et al.* (2020) Increase in the Number of Bone Marrow Osteoclast Precursors at Different Skeletal Sites, Particularly in Long Bone and Jaw Marrow in Mice Lacking IL-1RA. <u>Int J Mol Sci. 21 (11): 3774.</u>

38. Pluijmert, N.J. *et al.* (2021) Phosphorylcholine antibodies restrict infarct size and left ventricular remodelling by attenuating the unreperfused post-ischaemic inflammatory response. J Cell Mol Med. 25 (16): 7772-82.

39. Njock, M-K. (2022) Endothelial extracellular vesicles promote tumour growth by tumour-associated macrophage reprogramming <u>J Extracell Vesicles 2022</u> Jun;11(6):e12228.

Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.		
Guarantee	12 months from date of despatch		

Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA2389SBUV445 20471
Regulatory	For research purposes only
Related Produc Recommended Us	

MOUSE SEROBLOCK FcR (BUF041A) MOUSE SEROBLOCK FcR (BUF041B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-r	ad.com	Email: antibody_sales_uk@bio-r	ad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M395193:220321'

Printed on 12 Dec 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint