

Datasheet: MCA2393

BATCH NUMBER 161042

Description:	RAT ANTI MOUSE ER-HR3
Specificity:	ER-HR3
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	ER-HR3
Isotype:	IgG2c
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/50
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Adherent F1 (CBAxBL) bone marrow stromal cells.
RRID	AB_872053
Fusion Partners	Cells from immunised rats were fused with cells of the mouse P3 - X63 - Ag8.563 myeloma cell line.
Specificity	<p>Rat anti Mouse ER-HR3, clone ER-HR3 recognizes the murine antigen ER-HR3, a cell surface antigen expressed by Langerhans cells in epithelium, a subset of mature macrophages and dendritic cells located predominantly in haematopoietic and lymphoid organs. ER-HR3 demonstrates very low levels of expression on peripheral blood monocytes.</p> <p>During foetal development, ER-HR3 positive cells are localized to haematopoietic islands and are often associated with erythroid progenitor cells. ER-HR3 antigen may be involved in adult erythropoiesis and in the regulation of the immune response (de Jong et al. 1994).</p> <p>Rat anti Mouse ER-HR3, clone ER-HR3 does not inhibit antigen-specific T-cell proliferation.</p> <p>Rat anti Mouse ER-HR3, clone ER-HR3 recognizes two bands of ~69 kDa and a minor one of ~55 kDa under non-reducing conditions (de Jong et al. 1994).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> de Jong, J.P. <i>et al.</i> (1994) A monoclonal antibody (ER-HR3) against murine macrophages. I. Ontogeny, distribution and enzyme histochemical characterization of ER-HR3-positive cells. Cell Tissue Res. 275: 567-76. Oliveira, M.A. <i>et al.</i> (2003) Immature macrophages derived from mouse bone marrow produce large amounts of IL-12p40 after LPS stimulation. J Leukoc Biol. 74: 857-67. Throsby, M. <i>et al.</i> (2000) CD11c+ eosinophils in the murine thymus: developmental regulation and recruitment upon MHC class I-restricted thymocyte deletion. J Immunol. 165:1965-75. Grabbe, S. <i>et al.</i> (2002) Beta2 integrins are required for skin homing of primed T cells but not for priming naive T cells. J Clin Invest. 109: 183-92. Sonoda, Y. and Sasaki, K. (2012) Hepatic extramedullary hematopoiesis and macrophages in the adult mouse: histometrical and immunohistochemical studies. Cells Tissues Organs. 196: 555-64. Jacobsen, R.N. <i>et al.</i> (2014) Mobilization with granulocyte colony-stimulating factor blocks medullar erythropoiesis by depleting F4/80+VCAM1+CD169+ER-HR3+Ly6G+ erythroid island macrophages in the mouse. Exp Hematol. pii: S0301-472X(14)00139-8. Vogel, J. <i>et al.</i> (2003) Transgenic mice overexpressing erythropoietin adapt to

excessive erythrocytosis by regulating blood viscosity. [Blood. 102 \(6\): 2278-84.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2393>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rat IgG (STAR73...)	RPE
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight@550 , DyLight@650 , DyLight@800
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR16...)	DyLight@800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP
Goat Anti Rat IgG (STAR69...)	FITC

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M383921:210513'

Printed on 04 Sep 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)