

Datasheet: MCA2405A488

Description:	MOUSE ANTI HUMAN CD314:Alexa Fluor® 488
Specificity:	CD314
Other names:	NKG2D
Format:	ALEXA FLUOR® 488
Product Type:	Monoclonal Antibody
Clone:	1D11
Isotype:	IgG1
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Alexa Fluor® 488 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Alexa Fluor®488	495	519
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% sodium azide (NaN ₃)		
Stabilisers	1% bovine serum albumin		
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml		

Immunogen	NKL cells.
External Database Links	<p>UniProt: P26718 Related reagents</p> <p>Entrez Gene: 100528032 KLRC4-KLRK1 Related reagents</p>
Synonyms	D12S2489E, NKG2D
RRID	AB_2133270
Fusion Partners	Spleen cells from immunised RBF/DnJ mice were fused with cells of the p3 mouse myeloma cell line.
Specificity	<p>Mouse anti Human CD314 antibody, clone 1D11 recognizes CD314, also known as natural killer receptor G2 (NKG2D) and as killer cell lectin-like receptor subfamily K, member 1 (KLRK1).</p> <p>CD314 is a C-type lectin-like activating receptor which is expressed on most natural killer (NK) cells, CD8 T cells and gamma delta T cells. CD314 forms homodimers that signal through an associated DAP10 adaptor protein.</p> <p>Ligands of CD314 include MICA, MICB and UL16 binding protein (ULBP), which are inducibly expressed. Ligand binding to CD314 results in NK cell activation and potent co-stimulation of effector T cells.</p> <p>Mouse anti Human CD314 antibody, clone 1D11 is reported to inhibit T cell recognition of MICA (Bauer <i>et al.</i> 1999).</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> 1. Wu, J. <i>et al.</i> (2000) DAP10 and DAP12 form distinct, but functionally cooperative, receptor complexes in natural killer cells. J Exp Med. 192:1059-68. 2. Groh, V. <i>et al.</i> (2001) Costimulation of CD8alpha T cells by NKG2D via engagement by MIC induced on virus-infected cells. Nat Immunol. 2 (3): 255-60. 3. Roberts, A.I. <i>et al.</i> (2001) NKG2D receptors induced by IL-15 costimulate CD28-negative effector CTL in the tissue microenvironment. J Immunol. 167: 5527-30. 4. Gumperz, J. <i>et al.</i> (2002) Functionally distinct subsets of CD1d-restricted natural killer T cells revealed by CD1d tetramer staining. J Exp Med. 195:625-36. 5. Wu, J. <i>et al.</i> (2002) T cell antigen receptor engagement and specificity in the recognition of stress-inducible MHC class I-related chains by human epithelial gamma delta T cells. J Immunol. 169:1236-40. 6. Jinushi, M. <i>et al.</i> (2003) Autocrine/paracrine IL-15 that is required for type I IFN-mediated dendritic cell expression of MHC class I-related chain A and B is impaired in hepatitis C virus infection. J Immunol. 171 (10): 5423-9. 7. Groh, V. <i>et al.</i> (2003) Stimulation of T cell autoreactivity by anomalous expression of

- NKG2D and its MIC ligands in rheumatoid arthritis. [Proc Natl Acad Sci U S A. 100:9452-7](#)
8. Das, H. *et al.* (2004) Mechanisms of Vdelta1 gammadelta T cell activation by microbial components. [J Immunol. 172 \(11\): 6578-86.](#)
9. Holmen, C. *et al.* (2007) Anti endothelial cell autoantibodies selectively activate SAPK/JNK signalling in Wegener's granulomatosis. [J Am Soc Nephrol. 18: 2497-508.](#)
10. Sugita, J. *et al.* (2010) Differential effects of interleukin-12 and interleukin-15 on expansion of NK cell receptor-expressing CD8+ T cells. [Ann Hematol. 89: 115-20.](#)
11. Tanaka, J. *et al.* (2012) Expansion of NK cells from cord blood with antileukemic activity using GMP-compliant substances without feeder cells. [Leukemia. 26 \(5\): 1149-52.](#)
12. Matzner, P. *et al.* (2013) Resilience of the immune system in healthy young students to 30-hour sleep deprivation with psychological stress. [Neuroimmunomodulation. 20: 194-204.](#)
13. Voigt, J. *et al.* (2014) Human natural killer cells acting as phagocytes against *Candida albicans* and mounting an inflammatory response that modulates neutrophil antifungal activity. [J Infect Dis. 209 \(4\): 616-26.](#)
14. Tahrali, I. *et al.* (2019) CD3(-)CD56(+) NK cells display an inflammatory profile in RR-MS patients. [Immunol Lett. 216: 63-9.](#)

Further Reading 1. Groh, V. *et al.* (2003) Stimulation of T cell autoreactivity by anomalous expression of NKG2D and its MIC ligands in rheumatoid arthritis. [Proc Natl Acad Sci U S A. 100 \(16\): 9452-7.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Acknowledgements This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2405A488>
10041

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Alexa Fluor® 488 \(MCA928A488\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M415044:221215'

Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)