

Datasheet: MCA2405EL

#### **BATCH NUMBER 153124**

Description:	MOUSE ANTI HUMAN CD314:Low Endotoxin
Specificity:	CD314
Other names:	NKG2D
Format:	Low Endotoxin
Product Type:	Monoclonal Antibody
Clone:	1D11
Isotype:	lgG1
Quantity:	0.5 mg

# **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	<b>Not Determined</b>	Suggested Dilution
Flow Cytometry	•			1/50 - 1/200
Immunohistology - Frozen	•			
Immunohistology - Paraffin			•	
ELISA			•	
Immunoprecipitation	•			
Western Blotting			•	
Functional Assays				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein A fi supernatant	rom tissue culture
Buffer Solution	Phosphate buffered saline	
Preservative	None present	

# **Stabilisers Carrier Free** Yes **Endotoxin Level** < 0.01 EU/ug Approx. Protein IgG concentration 1.0 mg/ml Concentrations **Immunogen** NKL cells. **External Database UniProt:** Links P26718 Related reagents **Entrez Gene:** 100528032 KLRC4-KLRK1 Related reagents **Synonyms** D12S2489E, NKG2D **RRID** AB 609593 **Fusion Partners** Spleen cells from immunised RBF/DnJ mice were fused with cells of the p3 mouse myeloma cell line. **Specificity** Mouse anti Human CD314 antibody, clone 1D11 recognizes CD314, also known as natural killer receptor G2 (NKG2D) and as killer cell lectin-like receptor subfamily K, member 1 (KLRK1). CD314 is a C-type lectin-like activating receptor which is expressed on most natural killer (NK) cells, CD8 T cells and gamma delta T cells. CD314 forms homodimers that signal through an associated DAP10 adaptor protein. Ligands of CD314 include MICA, MICB and UL16 binding protein (ULBP), which are inducibly expressed. Ligand binding toCD314 results in NK cell activation and potent co-stimulation of effector T cells. Mouse anti Human CD314 antibody, clone 1D11 is reported to inhibit T cell recognition of MICA (Bauer et al. 1999). Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells in 100ul. Flow Cytometry References 1. Bauer, S. et al. (1999) Activation of NK cells and T cells by NKG2D, a receptor for

- stress-inducible MICA. Science. 285 (5428): 727-9.
- 2. Das, H. *et al.* (2004) Mechanisms of Vdelta1 gammadelta T cell activation by microbial components. <u>J Immunol. 172 (11): 6578-86.</u>
- 3. Groh, V. et al. (2001) Costimulation of CD8alphabeta T cells by NKG2D via engagement by MIC induced on virus-infected cells. <u>Nat Immunol. 2 (3): 255-60.</u>
- 4. Jinushi, M. et al. (2003) Autocrine/paracrine IL-15 that is required for type I

IFN-mediated dendritic cell expression of MHC class I-related chain A and B is impaired in hepatitis C virus infection. J Immunol. 171 (10): 5423-9. 5. Roberts, A.I. et al. (2001) NKG2D receptors induced by IL-15 costimulate CD28-negative effector CTL in the tissue microenvironment. J Immunol. 167: 5527-30. 6. Holmen, C. et al. (2007) Anti endothelial cell autoantibodies selectively activate SAPK/JNK signalling in Wegener's granulomatosis. J Am Soc Nephrol. 18: 2497-508. 7. Sugita, J. et al. (2010) Differential effects of interleukin-12 and interleukin-15 on expansion of NK cell receptor-expressing CD8+ T cells. Ann Hematol. 89: 115-20. 8. Gumperz, J. et al. (2002) Functionally distinct subsets of CD1d-restricted natural killer T cells revealed by CD1d tetramer staining. J Exp Med. 195:625-36. 9. Wu, J. et al. (2002) T cell antigen receptor engagement and specificity in the recognition of stress-inducible MHC class I-related chains by human epithelial gamma delta T cells. J Immunol. 169:1236-40. 10. Wu, J. et al. (2000) DAP10 and DAP12 form distinct, but functionally cooperative, receptor complexes in natural killer cells. <u>J Exp Med. 192:1059-68.</u> 11. Groh, V. et al. (2003) Stimulation of T cell autoreactivity by anomalous expression of NKG2D and its MIC ligands in rheumatoid arthritis. Proc Natl Acad Sci U S A. 100:9452-7 12. Voigt, J. et al. (2014) Human natural killer cells acting as phagocytes against Candida albicans and mounting an inflammatory response that modulates neutrophil antifungal activity. J Infect Dis. 209 (4): 616-26. 13. Matzner, P. et al. (2013) Resilience of the immune system in healthy young students to 30-hour sleep deprivation with psychological stress. Neuroimmunomodulation. 20: 14. Tahrali, I. et al. (2019) CD3-CD56+ NK cells display an inflammatory profile in RR-MS patients. Immunol Lett. Oct 04 [Epub ahead of print]. 15. Tanaka, J. et al. (2012) Expansion of NK cells from cord blood with antileukemic activity using GMP-compliant substances without feeder cells. Leukemia. 26 (5): 1149-52. **Further Reading** 1. Groh, V. et al. (2003) Stimulation of T cell autoreactivity by anomalous expression of NKG2D and its MIC ligands in rheumatoid arthritis. Proc Natl Acad Sci U S A. 100 (16): 9452-7. **Storage** Store at -20°C only. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10162 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2405EL">https://www.bio-rad-antibodies.com/SDS/MCA2405EL</a> 10162
Regulatory	For research purposes only

#### Related Products

#### **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12...)

Goat Anti Mouse IgG IgA IgM (STAR87...)

HRP

Goat Anti Mouse IgG (STAR76...)

Rabbit Anti Mouse IgG (STAR13...)

HRP

Goat Anti Mouse IgG (STAR70...)

FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) <u>FITC</u>
Goat Anti Mouse IgG (STAR77...) <u>HRP</u>

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

## **Recommended Negative Controls**

#### MOUSE IgG1 NEGATIVE CONTROL:Low Endotoxin (MCA928EL)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M366830:200529'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint