

## Datasheet: MCA2409GA

<b>Description:</b>	MOUSE ANTI HUMAN CD178
<b>Specificity:</b>	CD178
<b>Other names:</b>	FAS LIGAND
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	14C2
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			1/25 - 1/50
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation	▪			
Western Blotting		▪		

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) **Results maybe enhanced using membrane permeabilisation. Bio-Rad recommends the use of Leucoperm™ (Product code [BUF09](#)) prior to staining.**

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml

**External Database  
Links**

**UniProt:**

[P48023](#)   [Related reagents](#)

**Entrez Gene:**

[356](#)   FASLG   [Related reagents](#)

---

**Synonyms**

APT1LG1, CD95L, FASL, TNFSF6

---

**RRID**

AB\_566574

---

**Fusion Partners**

Spleen cells from immunised BALB/c mice were fused with cells of the P3U1 myeloma cell line.

---

**Specificity**

**Mouse anti Human CD178 antibody, clone 14C2** recognizes the human CD178, also known as Tumor necrosis factor ligand superfamily member 6, Fas ligand (FasL), Apoptosis antigen ligand or CD95 ligand. CD178 is a 281 amino acid, a ~40 kDa single pass type-II transmembrane glycoprotein bearing a single [intracellular FasL](#) domain and member of the tumor necrosis factor family .

CD178 is expressed by activated T lymphocytes and NK cells ([Leite-de-Moraes and Dy 1997](#)). The protein may exist as either a membrane bound or a cleaved soluble form ([Garcia et al. 2013](#)). CD178 plays an important role in T cell mediated cytotoxicity ([Jodo et al. 2005](#)). Binding of CD178 to Fas (CD95) results in the induction of apoptosis ([Ju et al. 1995](#)).

Mouse anti human CD178 antibody, clone 14C2 is reported to recognize a conformation dependent non-blocking epitope on CD178 ([Daburon et al. 2013](#)).

---

**Flow Cytometry**

Use 10ul of the suggested working dilution to label 1x10<sup>6</sup> cells in 100ul.

---

**References**

1. Legembre, P. *et al.* (2005) Amplification of Fas-mediated apoptosis in type II cells via microdomain recruitment. [Mol Cell Biol. 25 \(15\): 6811-20.](#)
2. Mesdaghi, M. *et al.* (2010) Natural killer cells in allergic rhinitis patients and nonatopic controls. [Int Arch Allergy Immunol. 153 \(3\): 234-8.](#)
3. Li, R. *et al.* (2014) Human heat shock protein-specific cytotoxic T lymphocytes display potent antitumour immunity in multiple myeloma. [Br J Haematol. 166 \(5\): 690-701.](#)
4. Ouwendijk, W.J. *et al.* (2014) Functional characterization of ocular-derived human alphaherpesvirus cross-reactive CD4 T cells. [J Immunol. 192: 3730-9.](#)
5. Matzner, P. *et al.* (2013) Resilience of the immune system in healthy young students to 30-hour sleep deprivation with psychological stress. [Neuroimmunomodulation. 20: 194-204.](#)
6. Sullivan, E.M. *et al.* (2014) NK cell genotype and phenotype at diagnosis of acute lymphoblastic leukemia correlate with postinduction residual disease. [Clin Cancer Res. 20 \(23\): 5986-94.](#)
7. Lindqvist CA *et al.* (2011) Both CD4+ FoxP3+ and CD4+ FoxP3- T cells from patients with B-cell malignancy express cytolytic markers and kill autologous leukaemic B cells *in vitro*. [Immunology. 133 \(3\): 296-306.](#)
8. Holmannova D *et al.* (2015) Inhibitory CD200R and proapoptotic CD95/CD95L molecules on innate immunity cells are modulated by cardiac surgery. [Perfusion. 30 \(7\): 543-55.](#)
9. Pachnio, A. *et al.* (2016) Cytomegalovirus Infection Leads to Development of High Frequencies of Cytotoxic Virus-Specific CD4+ T Cells Targeted to Vascular Endothelium. [PLoS Pathog. 12 \(9\): e1005832.](#)

---

**Storage**

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature

the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: 10040: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</a>
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">Alk. Phos.</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR8...)	<a href="#">DyLight@800</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight@488</a> , <a href="#">DyLight@680</a> , <a href="#">DyLight@800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

'M366842:200529'

Printed on 11 Aug 2020