

Datasheet: MCA2410PE

Description:	MOUSE ANTI MOUSE NOTCH 1:RPE		
Specificity:	NOTCH 1		
Format:	RPE		
Product Type:	Monoclonal Antibody		
Clone:	mN1A		
Isotype:	lgG1		
Quantity:	100 TESTS		

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further						
	information. For general protocol recommendations, please visit <u>www.bio-</u>						
	rad-antibodies.com/protocols. Yes No Not Determined Sugges						
	Flow Cytometry (1)	•			Neat		
	 Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls. (1) Membrane permeabilization is required for this application. The use of Leucoperm (Product Code <u>BUF09</u>) is recommended for this purpose. 						
Target Species	Mouse						
Species Cross Reactivity	Reacts with: Human N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.						
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized						
Reconstitution	Reconstitute with 1.0 ml distilled water Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.						
Max Ex/Em	Fluorophore RPE 488nm laser	Excitation Max 496	(nm) l	Emission Max (nm) 578			

Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃) 1% bovine serum albumin 5% sucrose
Immunogen	Synthetic peptide corresponding to cdc10-NCR region within mouse Notch1.
External Database Links	UniProt: Q01705 Related reagents Entrez Gene: 18128 Notch1 Related reagents
Synonyms	Motch
RRID	AB_609782
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse SP2/0 myeloma cell line.
Specificity	Mouse anti Mouse Notch 1 antibody, clone mN1A recognizes Notch 1, one of the four major transmembrane receptors (Notch 1-4) of the Notch signaling pathway, which is activated through binding to DSL domain-containing membrane-bound specific ligands. Notch 1 is expressed in a range of cells including haematopoietic cells in mouse foetal liver, adult thymus and bone marrow. Notch 1 signaling plays a role in follicular differentiation, tissue homeostasis, and in both CD4+ and CD8+ cell maturation in the thymus.
Specificity Flow Cytometry	major transmembrane receptors (Notch 1-4) of the Notch signaling pathway, which is activated through binding to DSL domain-containing membrane-bound specific ligands. Notch 1 is expressed in a range of cells including haematopoietic cells in mouse foetal liver, adult thymus and bone marrow. Notch 1 signaling plays a role in follicular differentiation, tissue homeostasis, and in both CD4+ and CD8+ cell maturation in the
	major transmembrane receptors (Notch 1-4) of the Notch signaling pathway, which is activated through binding to DSL domain-containing membrane-bound specific ligands. Notch 1 is expressed in a range of cells including haematopoietic cells in mouse foetal liver, adult thymus and bone marrow. Notch 1 signaling plays a role in follicular differentiation, tissue homeostasis, and in both CD4+ and CD8+ cell maturation in the thymus.
Flow Cytometry	 major transmembrane receptors (Notch 1-4) of the Notch signaling pathway, which is activated through binding to DSL domain-containing membrane-bound specific ligands. Notch 1 is expressed in a range of cells including haematopoietic cells in mouse foetal liver, adult thymus and bone marrow. Notch 1 signaling plays a role in follicular differentiation, tissue homeostasis, and in both CD4+ and CD8+ cell maturation in the thymus. Use 10µl of the suggested working dilution to label 1x10⁶ cells in 100µl 1. Ray, W.J. <i>et al.</i> (1999) Evidence for a physical interaction between presenilin and Notch. Proc Natl Acad Sci U S A. 96 (6): 3263-8. 2. Espinosa, L. <i>et al.</i> (2002) p65-NFkappaB synergizes with Notch to activate transcription by triggering cytoplasmic translocation of the nuclear receptor corepressor N-CoR. J Cell Sci. 115 (Pt 6): 1295-303. 3. Ren, M. and Cowell, J.K. (2011) Constitutive Notch pathway activation in murine ZMYM2-FGFR1-induced T-cell lymphomas associated with atypical myeloproliferative

	Prior to reconstitution store at +4°C.
	After reconstitution store at +4°C.
	DO NOT FREEZE.
	This product is photosensitive and should be protected from light.
Guarantee	12 months from date of despatch
Health And Safety	Material Safety Datasheet documentation #20487 available at:
Information	https://www.bio-rad-antibodies.com/SDS/MCA2410PE 20487
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

MOUSE SEROBLOCK FcR (BUF041A) MOUSE SEROBLOCK FcR (BUF041B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio	o-rad.com	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M440600:250523'

Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint