

Datasheet: MCA2411GA

Description:	MOUSE ANTI DOG CD34
Specificity:	CD34
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	1H6
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Dog
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Canine CD34 fusion protein.
External Database Links	<p>UniProt: Q28270 Related reagents</p> <p>Entrez Gene: 415130 CD34 Related reagents</p>
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS-1/FOX-NY myeloma cell line.
Specificity	<p>Mouse anti dog CD34 antibody, clone 1H6 recognizes the canine homologue of CD34, a glycosylated type 1 transmembrane protein of approximately 110 kDa (McSweeney et al. 1998) expressed on the cell surface of endothelial cells and haematopoietic stem cells.</p> <p>Mouse anti dog CD34 antibody, clone 1H6 is a key marker of canine hematopoietic progenitor cells and is reported for use in CD34+ enrichment assays, (Goerner et al. 2001) and (Horn et al. 2004).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
Western Blotting	MCA2411GA detects a band of approximately 110kDa.
References	<ol style="list-style-type: none"> Goerner, M. <i>et al.</i> (1999) The use of granulocyte colony-stimulating factor during retroviral transduction on fibronectin fragment CH-296 enhances gene transfer into hematopoietic repopulating cells in dogs. Blood. 94 (7): 2287-92. Bhattacharya, V. <i>et al.</i> (2000) Enhanced endothelialization and microvessel formation in polyester grafts seeded with CD34(+) bone marrow cells. Blood. 95 (2): 581-5. Goerner, M. <i>et al.</i> (2001) Sustained multilineage gene persistence and expression in dogs transplanted with CD34(+) marrow cells transduced by RD114-pseudotype oncoretrovirus vectors. Blood. 98 (7): 2065-70. Georges, G. <i>et al.</i> (2001) Engraftment of DLA-haploidentical marrow with ex vivo expanded, retrovirally transduced cytotoxic T lymphocytes. Blood. 98:3447-55. Horn, P.A. <i>et al.</i> (2004) Efficient lentiviral gene transfer to canine repopulating cells using an overnight transduction protocol. Blood. 103 (10): 3710-6. Avallone, G. <i>et al.</i> (2007) The spectrum of canine cutaneous perivascular wall tumors: morphologic, phenotypic and clinical characterization. Vet Pathol. 44 (5): 607-20. Palmieri, C. <i>et al.</i> (2013) Use of electron microscopy to classify canine perivascular wall tumors. Vet Pathol. 50 (2): 226-33. Bearden, R.N. <i>et al.</i> (2017) <i>In-vitro</i> characterization of canine multipotent stromal cells isolated from synovium, bone marrow, and adipose tissue: a donor-matched comparative study. Stem Cell Res Ther. 8 (1): 218. Trindade, A.B. <i>et al.</i> (2017) Mesenchymal-like stem cells in canine ovary show high differentiation potential. Cell Prolif. Oct 08 [Epub ahead of print].

10. Lee, S.H. *et al.* (2016) Impact of local injection of brain-derived neurotrophic factor-expressing mesenchymal stromal cells (MSCs) combined with intravenous MSC delivery in a canine model of chronic spinal cord injury. [Cytotherapy. Oct 28 \[Epub ahead of print\]](#).
11. Muir, P. *et al.* (2016) Autologous Bone Marrow-Derived Mesenchymal Stem Cells Modulate Molecular Markers of Inflammation in Dogs with Cruciate Ligament Rupture. [PLoS One. 11 \(8\): e0159095.](#)
12. Rajawat, Y.S. *et al.* (2021) *In Vivo* Gene Therapy for Canine SCID-X1 Using Cocal-Pseudotyped Lentiviral Vector. [Hum Gene Ther. 32 \(1-2\): 113-27.](#)
13. Grudzien, M. *et al.* (2021) A newly established canine NK-type cell line and its cytotoxic properties. [Vet Comp Oncol. 19 \(3\): 567-77.](#)
14. Tongu, E.A.O. *et al.* (2021) Allogenic mesenchymal stem cell-conditioned medium does not affect sperm parameters and mitigates early endometrial inflammatory responses in mares. [Theriogenology. 169: 1-8.](#)
15. Jaensch, S. *et al.* (2022) Clinicopathologic and immunophenotypic features in dogs with presumptive large granular lymphocyte leukaemia [Australian Veterinary Journal. \[Epub ahead of print\]](#).
16. Salari Sedigh, H. *et al.* (2023) *In vitro* investigation of canine periodontal ligament-derived mesenchymal stem cells: A possibility of promising tool for periodontal regeneration. [J Oral Biol Craniofac Res. 13 \(3\): 403-11.](#)
17. Papa, P.M. *et al.* (2023) Intratesticular transplantation of allogenic mesenchymal stem cells mitigates testicular destruction after induced heat stress in Miniature-horse stallions. [J Equine Vet Sci. 132: 104961.](#)
18. Rezaei, M. *et al.* (2019) Transplantation of Bone Marrow-Derived Mesenchymal Stem Cells, Platelet-Rich Plasma, and Fibrin Glue for Periodontal Regeneration. [Int J Periodontics Restorative Dent. 39 \(1\): e32-e45.](#)
19. Yang, V.K. *et al.* (2021) Intravenous administration of allogeneic Wharton jelly-derived mesenchymal stem cells for treatment of dogs with congestive heart failure secondary to myxomatous mitral valve disease. [Am J Vet Res. 82 \(6\): 487-93.](#)
20. Crain, S.K. *et al.* (2019) Extracellular Vesicles from Wharton's Jelly Mesenchymal Stem Cells Suppress CD4 Expressing T Cells Through Transforming Growth Factor Beta and Adenosine Signaling in a Canine Model. [Stem Cells Dev. 28 \(3\): 212-26.](#)
21. Sheng, R. *et al.* (2023) Prognostic significance of CD25 expression in dogs with a noninvasive diagnosis of B-cell lymphoma treated with CHOP chemotherapy. [Vet Comp Oncol. 21 \(1\): 28-35.](#)
22. Millanta, F. *et al.* (2020) Cytologic grading of canine and feline spindle-cell sarcomas of soft tissues and its correlation with histologic grading. [Top Companion Anim Med. 41: 100458.](#)

Further Reading

1. McSweeney, P. *et al.* (1996) Canine CD34: cloning of the cDNA and evaluation of an antiserum to recombinant protein. [Blood. 88:1992-2003.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2411GA 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight@488 , DyLight@550 , DyLight@650 , DyLight@680 , DyLight@800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M414900:221214'

Printed on 25 Mar 2024