

Datasheet: MCA2411PE

## **BATCH NUMBER 1705**

Description:	MOUSE ANTI DOG CD34:RPE
Specificity:	CD34
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	1H6
Isotype:	lgG1
Quantity:	100 TESTS

# **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Dog				
Product Form	Purified IgG conjug	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilize			
Reconstitution	Reconstitute with 1 ml distilled water				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)		
	RPE 488nm laser	496	578		
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue cultusupernatant				
Buffer Solution	Phosphate buffered saline				
Preservative	0.09% Sodium Azid	le			
Stabilisers 1% Bovine Serum Album	m Albumin				
	5% Sucrose				

Immunogen	Canine CD34 fusion protein.
External Database Links	UniProt:  Q28270 Related reagents
	Entrez Gene:  415130 CD34 Related reagents
RRID	AB_609594
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS-1/FOX-NY myeloma cell line.
Specificity	<b>Mouse anti dog CD34 antibody, clone 1H6</b> recognizes the canine homologue of CD34, a glycosylated type 1 transmembrane protein of approximately 110 kDa ( <u>McSweeney et al. 1998</u> ) expressed on the cell suface of endothelial cells and haematopoietic stem cells.
	Mouse anti dog CD34 antibody, clone 1H6 is a key marker of canine hematopoietic progenitor cells and is reported for use in CD34+ enrichment assays, ( <u>Goerner et al. 2001</u> ) and ( <u>Horn et al. 2004</u> ).
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.
References	<ol> <li>Goerner, M. <i>et al.</i> (1999) The use of granulocyte colony-stimulating factor during retroviral transduction on fibronectin fragment CH-296 enhances gene transfer into hematopoietic repopulating cells in dogs. <u>Blood. 94 (7): 2287-92.</u></li> <li>Bhattacharya, V. <i>et al.</i> (2000) Enhanced endothelialization and microvessel formation in polyester grafts seeded with CD34(+) bone marrow cells. <u>Blood. 95 (2): 581-5.</u></li> <li>Goerner, M. <i>et al.</i> (2001) Sustained multilineage gene persistence and expression in dogs transplanted with CD34(+) marrow cells transduced by RD114-pseudotype oncoretrovirus vectors. <u>Blood. 98 (7): 2065-70.</u></li> <li>Georges, G. <i>et al.</i> (2001) Engraftment of DLA-haploidentical marrow with ex vivo expanded, retrovirally transduced cytotoxic T lymphocytes. <u>Blood. 98:3447-55.</u></li> <li>Horn, P.A. <i>et al.</i> (2004) Efficient lentiviral gene transfer to canine repopulating cells</li> </ol>

- 5. Horn, P.A. *et al.* (2004) Efficient lentiviral gene transfer to canine repopulating cells using an overnight transduction protocol. <u>Blood</u>. 103 (10): 3710-6.
- 6. Avallone, G. *et al.* (2007) The spectrum of canine cutaneous perivascular wall tumors: morphologic, phenotypic and clinical characterization. <u>Vet Pathol. 44 (5): 607-20.</u>
- 7. Palmieri, C. *et al.* (2013) Use of electron microscopy to classify canine perivascular wall tumors. <u>Vet Pathol. 50 (2): 226-33.</u>
- 8. Bearden, R.N. *et al.* (2017) *In-vitro* characterization of canine multipotent stromal cells isolated from synovium, bone marrow, and adipose tissue: a donor-matched comparative study. <u>Stem Cell Res Ther. 8 (1): 218.</u>
- 9. Trindade, A.B. *et al.* (2017) Mesenchymal-like stem cells in canine ovary show high differentiation potential. <u>Cell Prolif. Oct 08 [Epub ahead of print].</u>
- 10. Lee, S.H. *et al.* (2016) Impact of local injection of brain-derived neurotrophic factor-expressing mesenchymal stromal cells (MSCs) combined with intravenous MSC delivery in a canine model of chronic spinal cord injury. Cytotherapy. Oct 28 [Epub ahead of print].

- 11. Muir, P. *et al.* (2016) Autologous Bone Marrow-Derived Mesenchymal Stem Cells Modulate Molecular Markers of Inflammation in Dogs with Cruciate Ligament Rupture. PLoS One. 11 (8): e0159095.
- 12. Rajawat, Y.S. *et al.* (2021) *In Vivo* Gene Therapy for Canine SCID-X1 Using Cocal-Pseudotyped Lentiviral Vector. <u>Hum Gene Ther. 32 (1-2): 113-27.</u>
- 13. Grudzien, M. *et al.* (2021) A newly established canine NK-type cell line and its cytotoxic properties. <u>Vet Comp Oncol. 19 (3): 567-77.</u>
- 14. Tongu, E.A.O. *et al.* (2021) Allogenic mesenchymal stem cell-conditioned medium does not affect sperm parameters and mitigates early endometrial inflammatory responses in mares. <u>Theriogenology</u>. 169: 1-8.
- 15. Jaensch, S. *et al.* (2022) Clinicopathologic and immunophenotypic features in dogs with presumptive large granular lymphocyte leukaemia <u>Australian Veterinary Journal.</u> [Epub ahead of print].

Further Reading	1. McSweeney, P. <i>et al.</i> (1996) Canine CD34: cloning of the cDNA and evaluation of a antiserum to recombinant protein. <u>Blood. 88:1992-2003.</u>
Storage	Prior to reconstitution store at +4°C.
	After reconstitution store at +4°C.
	DO NOT FREEZE. This product is photosensitive and should be protected from light.
	Should this product contain a precipitate we recommend microcentrifugation before us
Guarantee	12 months from date of despatch
Health And Safety	Material Safety Datasheet documentation #20487 available at:
Information	https://www.bio-rad-antibodies.com/SDS/MCA2411PE
	20487
Regulatory	For research purposes only

## Related Products

America

### **Recommended Negative Controls**

#### MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA928PE)

North & South Tel: +1 800 265 7376 Worldwide Tel: +44 (0)1865 852 70

Fax: +1 919 878 3751 F
Email: antibody\_sales\_us@bio-rad.com E

Fax: +49 (0) 89 8090 95 50

Tel: +49 (0) 89 8090 95 21

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M375482:210104'

# Printed on 25 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint