

## Datasheet: MCA2413GA

**BATCH NUMBER 161355**

<b>Description:</b>	MOUSE ANTI CHICKEN CD45
<b>Specificity:</b>	CD45
<b>Other names:</b>	LEUCOCYTE COMMON ANTIGEN
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	UM16-6
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/100 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Chicken
<b>Species Cross Reactivity</b>	Does not react with: Turkey
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline

<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	COS cells transfected with CD45 gene.
<b>Specificity</b>	<p><b>Mouse anti Chicken CD45 antibody, clone UM16-6</b> recognizes chicken CD45, also known as leucocyte common antigen. CD45 is a heavily-glycosylated transmembrane protein tyrosine phosphatase (PTPase) expressed by all nucleated cells of haematopoietic origin. Variation in the expression of a particular CD45 isoform, is regulated during the haematopoietic development of the different cell lineages.</p> <p>CD45 is essential for antigen-induced signal transduction through the antigen receptor and as with other PTPase family members, acts in balance with protein tyrosine kinases, causing the dephosphorylation of negative regulatory tyrosine sites. De-phosphorylation by CD45, is required for the activation of the src-family kinases p56<sup>lck</sup> and p59<sup>fyn</sup>.</p> <p>Chicken CD45 has an additional cysteine residue near the transmembrane region compared to human and shark CD45 (<a href="#">Okumura et al. 1996</a>). The overall domain structure between mammalian and chicken CD45 appears to be conserved, but the sequence homology between the extracellular regions is very low.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>Histology Positive Control Tissue</b>	Chicken lung
<b>References</b>	<ol style="list-style-type: none"> <li>1. Pavlova, S. <i>et al.</i> (2010) <i>In vitro</i> and <i>in vivo</i> characterization of glycoprotein C-deleted infectious laryngotracheitis virus. <a href="#">J Gen Virol. 91:847-57.</a></li> <li>2. Watrang, E. <i>et al.</i> (2015) CD107a as a marker of activation in chicken cytotoxic T cells. <a href="#">J Immunol Methods. 419: 35-47.</a></li> <li>3. Reddy SK <i>et al.</i> (2008) The BAFF-Interacting receptors of chickens. <a href="#">Dev Comp Immunol. 32 (9): 1076-87.</a></li> <li>4. Ulrich-Lynge SL <i>et al.</i> (2015) The consequence of low mannose-binding lectin plasma concentration in relation to susceptibility to <i>Salmonella infantis</i> in chickens. <a href="#">Vet Immunol Immunopathol. 163 (1-2): 23-32.</a></li> <li>5. Eren, U. <i>et al.</i> (2016) The several elements of intestinal innate immune system at the beginning of the life of broiler chicks. <a href="#">Microsc Res Tech. 79 (7): 604-14.</a></li> <li>6. Röhe, I. <i>et al.</i> (2017) Effect of feeding soybean meal and differently processed peas on the gut mucosal immune system of broilers <a href="#">Poultry Science. Feb 23 [Epub ahead of print]</a></li> <li>7. Czerwiński, J. <i>et al.</i> (2015) The use of genetically modified Roundup Ready soyabean meal and genetically modified MON 810 maize in broiler chicken diets. Part 1. Effects on performance and blood lymphocyte subpopulations <a href="#">J Anim Feed Sci 24: 134-43.</a></li> <li>8. Kjærup, R.B. <i>et al.</i> (2017) Comparison of growth performance and immune parameters of three commercial chicken lines used in organic production. <a href="#">Vet Immunol Immunopathol. 187: 69-79.</a></li> </ol>

9. Larsen, F.T. *et al.* (2019) Immunoprofiling of peripheral blood from infectious bronchitis virus vaccinated MHC-B chicken lines - Monocyte MHC-II expression as a potential correlate of protection. [Dev Comp Immunol. 96: 93-102.](#)
10. Naghizadeh, M. *et al.* (2019) Rapid whole blood assay using flow cytometry for measuring phagocytic activity of chicken leukocytes. [Vet Immunol Immunopathol. 207: 53-61.](#)
11. Fenzl, L. *et al.* (2017)  $\gamma\delta$  T cells represent a major spontaneously cytotoxic cell population in the chicken. [Dev Comp Immunol. 73: 175-83.](#)
12. Harvanová, D. *et al.* (2014) Isolation, cultivation and characterisation of pigeon osteoblasts seeded on xenogeneic demineralised cancellous bone scaffold for bone grafting. [Vet Res Commun. 38 \(3\): 221-8.](#)

---

<b>Further Reading</b>	<ol style="list-style-type: none"> <li>1. Symons, A. <i>et al.</i> (1999) Domain organization of the extracellular region of CD45. <a href="#">Protein Eng. 12 (10): 885-92.</a></li> <li>2. Okumura, M. <i>et al.</i> (1996) Comparison of CD45 extracellular domain sequences from divergent vertebrate species suggests the conservation of three fibronectin type III domains. <a href="#">J Immunol. 157 (4): 1569-75.</a></li> <li>3. Jung, E.J. <i>et al.</i> (1997) Phosphorylation of chicken protein tyrosine phosphatase 1 by casein kinase II <i>in vitro</i>. <a href="#">Experimental and Molecular Medicine 29(4): 229-33.</a></li> </ol>
------------------------	--

---

<b>Storage</b>	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p>
----------------	--

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	<p>Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2413GA">https://www.bio-rad-antibodies.com/SDS/MCA2413GA</a> 10040</p>
--------------------------------------	---

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR8...)	<a href="#">DyLight@800</a>
Human Anti Mouse IgG2a (HCA037...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">Alk. Phos.</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight@488</a> , <a href="#">DyLight@550</a> , <a href="#">DyLight@650</a> , <a href="#">DyLight@680</a> , <a href="#">DyLight@800</a> ,

	<a href="#">FITC</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M383824:210513'

**Printed on 25 Mar 2023**

---

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)