

Datasheet: MCA2413GA BATCH NUMBER 161355

Description:	MOUSE ANTI CHICKEN CD45
Specificity:	CD45
Other names:	LEUCOCYTE COMMON ANTIGEN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	UM16-6
Isotype:	lgG2a
Quantity:	0.1 mg

Product Details

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				1/100 - 1/200
Immunohistology - Frozen				
Immunohistology - Paraffin			•	
ELISA			-	
Immunoprecipitation	-			
Western Blotting			•	
Where this antibody has a necessarily exclude its us a guide only. It is recommon system using appropriate	e in such iended th	n procedu nat the us	res. Suggested workir er titrates the antibody	g dilutions are given a

Target Species	Chicken	
Species Cross Reactivity	Does not react with:Turkey	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant	
Buffer Solution	Phosphate buffered saline	

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)			
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml			
Immunogen	COS cells transfected with CD45 gene.			
Specificity	Mouse anti Chicken CD45 antibody, clone UM16-6 recognizes chicken CD45, also known as leucocyte common antigen. CD45 is a heavily-glycosylated transmembrane protein tyrosine phosphatase (PTPase) expressed by all nucleated cells of haematopoietic origin. Variation in the expression of a particular CD45 isoform, is regulated during the haematopoietic development of the different cell lineages.			
	CD45 is essential for antigen-induced signal transduction through the antigen receptor and as with other PTPase family members, acts in balance with protein tyrosine kinases, causing the dephosphorylation of negative regulatory tyrosine sites. De-phosphorylation by CD45, is required for the activation of the src-family kinases p56 ^{lck} and p59 ^{fyn} .			
	Chicken CD45 has an additional cysteine residue near the transmembrane region compared to human and shark CD45 (<u>Okumura <i>et al.</i> 1996</u>). The overall domain structure between mammalian and chicken CD45 appears to be conserved, but the sequence homology between the extracellular regions is very low.			
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.			
Histology Positive Control Tissue	Chicken lung			
References	 Reddy SK <i>et al.</i> (2008) The BAFF-Interacting receptors of chickens. <u>Dev Comp Immunol. 32 (9): 1076-87.</u> Pavlova, S. <i>et al.</i> (2010) <i>In vitro</i> and <i>in vivo</i> characterization of glycoprotein C-deleted infectious laryngotracheitis virus. <u>J Gen Virol. 91:847-57.</u> Harvanová, D. <i>et al.</i> (2014) Isolation, cultivation and characterisation of pigeon osteoblasts seeded on xenogeneic demineralised cancellous bone scaffold for bone grafting. <u>Vet Res Commun. 38 (3): 221-8.</u> Ulrich-Lynge SL <i>et al.</i> (2015) The consequence of low mannose-binding lectin plasma concentration in relation to susceptibility to <i>Salmonella infantis</i> in chickens. <u>Vet Immunol Immunopathol. 163 (1-2): 23-32.</u> Wattrang, E. <i>et al.</i> (2015) CD107a as a marker of activation in chicken cytotoxic T cells. <u>J Immunol Methods. 419: 35-47.</u> Czerwiński, J. <i>et al.</i> (2015) The use of genetically modified Roundup Ready soyabean meal and genetically modified MON 810 maize in broiler chicken diets. Part 1. Effects on performance and blood lymphocyte subpopulations <u>J Anim Feed Sci 24: 134-43.</u> Eren, U. <i>et al.</i> (2017) Effect of feeding soybean meal and differently processed peas on the gut mucosal immune system of broilers <u>Poultry Science. Feb 23 [Epub ahead of print]</u> 			

	 9. Kjærup, R.B. <i>et al.</i> (2017) Comparison of growth performance of three commercial chicken lines used in organic production. <u>187: 69-79.</u> 10. Fenzl, L. <i>et al.</i> (2017) γδ T cells represent a major spontane population in the chicken. <u>Dev Comp Immunol. 73: 175-83.</u> 11. Larsen, F.T. <i>et al.</i> (2019) Immunoprofiling of peripheral blood virus vaccinated MHC-B chicken lines - Monocyte MHC-II expression of protection. <u>Dev Comp Immunol. 96: 93-102.</u> 12. Naghizadeh, M. <i>et al.</i> (2019) Rapid whole blood assay usin measuring phagocytic activity of chicken leukocytes. <u>Vet Immunot 53-61.</u> 13. Wattrang, E. <i>et al.</i> (2020) Immune responses upon experime <i>rhusiopathiae</i> infection of naïve and vaccinated chickens. <u>Vet F</u> 14. Alber, A. <i>et al.</i> (2019) Avian Pathogenic Escherichia coli (All Immunomodulation of Respiratory Granulocytes and Mononucle Reporter Transgenic Chickens. <u>Front Immunol. 10: 3055.</u> 	Vet Immunol Immunopathol. eously cytotoxic cell d from infectious bronchitis ession as a potential g flow cytometry for hol Immunopathol. 207: mental Erysipelothrix Res. 51 (1): 114. PEC) Strain-Dependent
Further Reading	 Okumura, M. <i>et al.</i> (1996) Comparison of CD45 extracellular divergent vertebrate species suggests the conservation of three domains. <u>J Immunol. 157 (4): 1569-75.</u> Jung, E.J. <i>et al.</i> (1997) Phosphorylation of chicken protein ty casein kinase II <i>in vitro</i>. <u>Experimental and Molecular Medicine 2</u> Symons, A. <i>et al.</i> (1999) Domain organization of the extracel <u>Protein Eng. 12 (10): 885-92.</u> 	e fibronectin type III rosine phosphatase 1 by 29(4): 229-33.
Storage	This product is shipped at ambient temperature. It is recommer -20°C on receipt. When thawed, aliquot the sample as needed. short term use (up to 4 weeks) and store the remaining aliquots Avoid repeated freezing and thawing as this may denature the frost-free freezers is not recommended.	Keep aliquots at 2-8°C for s at -20°C.
Guarantee	12 months from date of despatch	
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2413GA 10040	
Regulatory	For research purposes only	

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12)	<u>RPE</u>
Goat Anti Mouse IgG IgA IgM (STAR87	.) <u>HRP</u>
Goat Anti Mouse IgG (STAR76)	<u>RPE</u>
Rabbit Anti Mouse IgG (STAR13)	<u>HRP</u>

Goat Ant	i Mouse IgG (STAR70)	<u>FIT</u>	<u>C</u>			
Goat Anti Mouse IgG (H/L) (STAR117)			Alk. Phos., DyLight®488, DyLight®550,			
		Dyl	<u>ight®650</u> , <u>DyLight®680</u>), DyLight®80	<u>00,</u>	
		FIT	<u>C, HRP</u>			
Rabbit A	nti Mouse IgG (STAR9)	<u>FIT</u>	<u>C</u>			
Goat Anti Mouse IgG (STAR77)		HR	HRP			
Goat Ant	i Mouse IgG (Fc) (STAR120) <u>FIT</u>	<u>C, HRP</u>			
North & South America	Tel: +1 800 265 7376 Work Fax: +1 919 878 3751	dwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50	

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M383824:210513'

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint

Email: antibody_sales_us@bio-rad.com