

Datasheet: MCA2420T

Description:	MOUSE ANTI HUMAN CD62P
Specificity:	CD62P
Other names:	P-SELECTIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	Psel.KO.2.12
Isotype:	IgG1
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Reacts with: Goat, Rat, Sheep N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml

Immunogen	CD62P transfected 300.19 cells.
External Database Links	<p>UniProt: P16109 Related reagents</p> <p>Entrez Gene: 6403 SELP Related reagents</p>
Synonyms	GMRP, GRMP
Fusion Partners	Spleen cells from immunised CD62P knock-out mice (Strain C57/B6) were fused with cells of the NS-1 myeloma cell line.
Specificity	<p>Mouse anti Human CD62P antibody, clone Psel.KO.2.12 recognizes the CD62P cell surface antigen, a ~140 kDa glycoprotein also known as P-selectin.</p> <p>CD62P is expressed by activated platelets and endothelial cells, and plays an important role in adhesive processes between leucocytes and endothelial cells.</p> <p>Mouse anti Human CD62P antibody, clone Psel.KO.2.12 inhibits P-selectin-dependent adhesion between HL60 cells and P-selectin transfected COS cells (Massaquer et al. 2000).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Massaquer, A. <i>et al.</i> (2000) Production and characterization of monoclonal antibodies against conserved epitopes of P-selectin (CD62P). Tissue Antigens. 56 (2): 117-28. 2. Massaquer, A. <i>et al.</i> (2003) Characterization of platelet and soluble-porcine P-selectin (CD62P). Vet Immunol Immunopathol. 96 (3-4): 169-81. 3. Massaquer, A. <i>et al.</i> (2002) Reactivity of CD62P workshop mAbs with resting and activated platelets from different animal species. In: Leucocyte Typing VII. Edited by Mason, D. <i>et al.</i> Oxford University Press, pp 342-3. 4. Major, T.C. <i>et al.</i> (2010) The attenuation of platelet and monocyte activation in a rabbit model of extracorporeal circulation by a nitric oxide releasing polymer. Biomaterials. 31: 2736-45. 5. Johnson, C.A. Jr. <i>et al.</i> (2008) Flow cytometric assays for quantifying activated ovine platelets. Artif Organs. 32: 136-45. 6. Johnson, C.A. Jr. <i>et al.</i> (2011) Platelet activation in ovinos undergoing sham surgery or implant of the second generation PediaFlow pediatric ventricular assist device. Artif Organs. 35 (6): 602-13. 7. Johnson, C.A. Jr. <i>et al.</i> (2011) Biocompatibility assessment of the first generation PediaFlow pediatric ventricular assist device. Artif Organs. 35 (1): 9-21. 8. Dasse, K.A. <i>et al.</i> (2007) Assessment of hydraulic performance and biocompatibility of a MagLev centrifugal pump system designed for pediatric cardiac or cardiopulmonary support. ASAIO J. 53 (6): 771-7. 9. Ding, J. <i>et al.</i> (2015) Quantification of Shear-Induced Platelet Activation: High Shear Stresses for Short Exposure Time. Artif Organs. 39 (7): 576-83. 10. Tran GT <i>et al.</i> (2010) Membrane attack complex of complement is not essential for immune mediated demyelination in experimental autoimmune neuritis. J Neuroimmunol. 229 (1-2): 98-106. 11. Foruzanmehr, M. <i>et al.</i> (2014) Nano-structure TiO₂ film coating on 316L stainless steel via sol-gel technique for blood compatibility improvement. Nanomedicine Journal 1 (3): 128-36.
Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>Storage in frost-free freezers is not recommended.</p> <p>This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature</p>

the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life	18 months from date of despatch.
-------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Rabbit Anti Mouse IgG (STAR8...)	DyLight@800
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight@488 , DyLight@549 , DyLight@649 , DyLight@680 , DyLight@800 , FITC , HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Human Anti Mouse IgG1 (HCA036...)	HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M323834:180727'

Printed on 18 Sep 2018