

Datasheet: MCA2430F

Description:	MOUSE ANTI BOVINE CD25:FITC
Specificity:	CD25
Other names:	IL-2R ALPHA CHAIN
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	IL-A111
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/5

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Bovine

Species Cross Reactivity

Reacts with: Sheep

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
FITC	490	525

Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
External Database Links	<p>UniProt: P12342 Related reagents</p> <p>Entrez Gene: 281861 IL2RA Related reagents</p>
RRID	AB_2249049
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the X63.Ag8.653 myeloma cell line.
Specificity	<p>Mouse anti Bovine CD25 antibody, clone IL-A111 recognizes the bovine CD25 cell surface antigen, a ~55 kDa glycoprotein also known as Interleukin-2 receptor alpha chain. Bovine CD25 is expressed by activated T cells.</p> <p>Mouse anti Bovine CD25 antibody, clone IL-A111 is reported to block the IL-2 driven proliferation of Con A-induced blast cells/ bovine lymphocytes (Naessens et al. 1992).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul
References	<ol style="list-style-type: none"> Naessens, J. <i>et al.</i> (1992) Selection of BoCD25 monoclonal antibodies by screening mouse L cells transfected with the bovine p55-interleukin-2 (IL-2) receptor gene. Immunology. 76 (2): 305-9. Howard, C.J. & Naessens, J. (1993) Summary of workshop findings for cattle (tables 1 and 2). Vet Immunol Immunopathol. 39 (1-3): 25-47. Howard, C.J. <i>et al.</i> (1997) Identification of two distinct populations of dendritic cells in afferent lymph that vary in their ability to stimulate T cells. J Immunol. 159 (11): 5372-82. Naessens, J. <i>et al.</i> (1993) Cross-reactivity of workshop antibodies with cells from domestic and wild ruminants. Vet Immunol Immunopathol. 39 (1-3): 283-90. Evans, C.W. <i>et al.</i> (1994) Antigen recognition and activation of ovine gamma delta T cells. Immunology. 82 (2): 229-37. Campbell, J.D. <i>et al.</i> (1998) A novel cell surface proliferation-associated marker expressed on T cells and up-regulated on germinal center B cells. J Leukoc Biol. 63 (5): 567-74. Connelley, T. <i>et al.</i> (2011) NKp46 defines ovine cells that have characteristics corresponding to NK cells. Vet Res. 42: 37. Menge, C. <i>et al.</i> (2004) Phenotypic and functional characterization of intraepithelial lymphocytes in a bovine ligated intestinal loop model of enterohaemorrhagic Escherichia coli infection. J Med Microbiol. 53: 573-9. Rhodes, S.G. <i>et al.</i> (1999) Differential cytokine responses of CD4+ and CD8+ T cells in response to bovine viral diarrhoea virus in cattle. J Gen Virol. 80 : 1673-9.

10. Piper, E.K. *et al.* (2009) Immunological profiles of *Bos taurus* and *Bos indicus* cattle infested with the cattle tick, *Rhipicephalus (Boophilus) microplus*. [Clin Vaccine Immunol. 16: 1074-86.](#)
11. Woolums, A.R. *et al.* (2013) Effect of calf age and administration route of initial multivalent modified-live virus vaccine on humoral and cell-mediated immune responses following subsequent administration of a booster vaccination at weaning in beef calves. [Am J Vet Res. 74: 343-54.](#)
12. McInnes, E. *et al.* (1999) Phenotypic analysis of local cellular responses in calves infected with bovine respiratory syncytial virus. [Immunology. 96 \(3\): 396-403.](#)
13. Maślanka, T. *et al.* (2012) The presence of CD25 on bovine WC1+ $\gamma\delta$ T cells is positively correlated with their production of IL-10 and TGF- β , but not IFN- γ . [Pol J Vet Sci. 15: 11-20.](#)
14. Menge, C. *et al.* (2003) Verotoxin 1 from *Escherichia coli* affects Gb3/CD77+ bovine lymphocytes independent of interleukin-2, tumor necrosis factor-alpha, and interferon-alpha. [Exp Biol Med \(Maywood\). 228: 377-86.](#)
15. Menge, C. *et al.* (1999) Shiga toxin 1 from *Escherichia coli* blocks activation and proliferation of bovine lymphocyte subpopulations *in vitro*. [Infect Immun. 67: 2209-17.](#)
16. Constantinoiu, C.C. *et al.* (2010) Local immune response against larvae of *Rhipicephalus (Boophilus) microplus* in *Bos taurus indicus* and *Bos taurus taurus* cattle. [Int J Parasitol. 40: 865-75.](#)
17. Maślanska, T. and Jaroszewski, J.J. (2013) In vitro effects of meloxicam on the number, Foxp3 expression, production of selected cytokines, and apoptosis of bovine CD25+CD4+ and CD25-CD4+ cells. [J Vet Sci. 14 \(2\): 125-34.](#)
18. Brodzki, P. *et al.* (2014) Phenotyping of leukocytes and granulocyte and monocyte phagocytic activity in the peripheral blood and uterus of cows with endometritis. [Theriogenology. 82 \(3\): 403-10.](#)
19. Zoldan, K. *et al.* (2014) Increase of CD25 expression on bovine neutrophils correlates with disease severity in post-partum and early lactating dairy cows. [Dev Comp Immunol. 47 \(2\): 254-63.](#)
20. Kang, H.J. *et al.* (2016) Effects of Ambient Temperature on Growth Performance, Blood Metabolites, and Immune Cell Populations in Korean Cattle Steers. [Asian-Australas J Anim Sci. 29 \(3\): 436-43.](#)
21. Schmidt, N. *et al.* (2018) Decreased STEC shedding by cattle following passive and active vaccination based on recombinant *Escherichia coli* Shiga toxoids. [Vet Res. 49 \(1\): 28.](#)
22. Higgins, J.L. *et al.* (2018) Cell mediated immune response in goats after experimental challenge with the virulent *Brucella melitensis* strain 16M and the reduced virulence strain Rev. 1. [Vet Immunol Immunopathol. 202: 74-84.](#)
23. Brodzki, P. *et al.* (2019) Selected leukocyte subpopulations in peripheral blood and uterine washings in cows before and after intrauterine administration of cefapirin and methisoprinol. [Anim Sci J. Nov 06 \[Epub ahead of print\].](#)
24. Rivalde, M.A. *et al.* (2020) BVDV permissiveness and lack of expression of co-stimulatory molecules on PBMCs from calves pre-infected with BVDV. [Comp Immunol Microbiol Infect Dis. 68: 101388.](#)
25. de Araújo, F.F. *et al.* (2019) Distinct immune response profile during *Rhipicephalus (Boophilus) microplus* infestations of guzerat dairy herd according to the maternal lineage ancestry (mitochondrial DNA). [Vet Parasitol. 273: 36-44.](#)

26. Colombatti Olivieri, M.A. *et al.* (2021) Evaluation of a virulent strain of *Mycobacterium avium* subsp. *paratuberculosis*. used as a heat-killed vaccine. [Vaccine. Nov 10;S0264-410X\(21\)01433-X.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M385393:210513'

Printed on 16 Nov 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)