

Datasheet: MCA2430GA

BATCH NUMBER 157704

Description:	MOUSE ANTI BOVINE CD25
Specificity:	CD25
Other names:	IL-2R ALPHA CHAIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	IL-A111
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Bovine
Species Cross Reactivity	<p>Reacts with: Sheep</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture

supernatant

Buffer Solution Phosphate buffered saline

Preservative Stabilisers 0.09% Sodium Azide

Carrier Free Yes

Approx. Protein Concentrations IgG concentration 1.0 mg/ml

External Database Links

UniProt:

[P12342](#) [Related reagents](#)

Entrez Gene:

[281861](#) IL2RA [Related reagents](#)

Fusion Partners Spleen cells from immunised BALB/c mice were fused with cells of the X63.Ag8.653 myeloma cell line.

Specificity **Mouse anti Bovine CD25 antibody, clone IL-A111** recognizes the bovine CD25 cell surface antigen, a ~55 kDa glycoprotein also known as Interleukin-2 receptor alpha chain. Bovine CD25 is expressed by activated T cells.

Mouse anti Bovine CD25 antibody, clone IL-A111 is reported to block the IL-2 driven proliferation of Con A-induced blast cells/ bovine lymphocytes ([Naessens et al. 1992](#)).

Flow Cytometry Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul

References

1. Naessens, J. *et al.* (1992) Selection of BoCD25 monoclonal antibodies by screening mouse L cells transfected with the bovine p55-interleukin-2 (IL-2) receptor gene. [Immunology. 76 \(2\): 305-9.](#)
2. Howard, C.J. & Naessens, J. (1993) Summary of workshop findings for cattle (tables 1 and 2). [Vet Immunol Immunopathol. 39 \(1-3\): 25-47.](#)
3. Howard, C.J. *et al.* (1997) Identification of two distinct populations of dendritic cells in afferent lymph that vary in their ability to stimulate T cells. [J Immunol. 159 \(11\): 5372-82.](#)
4. Naessens, J. *et al.* (1993) Cross-reactivity of workshop antibodies with cells from domestic and wild ruminants. [Vet Immunol Immunopathol. 39 \(1-3\): 283-90.](#)
5. Evans, C.W. *et al.* (1994) Antigen recognition and activation of ovine gamma delta T cells. [Immunology. 82 \(2\): 229-37.](#)
6. Campbell, J.D. *et al.* (1998) A novel cell surface proliferation-associated marker expressed on T cells and up-regulated on germinal center B cells. [J Leukoc Biol. 63 \(5\): 567-74.](#)
7. Connelley, T. *et al.* (2011) NKp46 defines ovine cells that have characteristics corresponding to NK cells. [Vet Res. 42: 37.](#)
8. Menge, C. *et al.* (2004) Phenotypic and functional characterization of intraepithelial lymphocytes in a bovine ligated intestinal loop model of enterohaemorrhagic Escherichia

- coli infection. [J Med Microbiol. 53: 573-9.](#)
9. Rhodes, S.G. *et al.* (1999) Differential cytokine responses of CD4+ and CD8+ T cells in response to bovine viral diarrhoea virus in cattle. [J Gen Virol. 80 : 1673-9.](#)
10. Piper, E.K. *et al.* (2009) Immunological profiles of *Bos taurus* and *Bos indicus* cattle infested with the cattle tick, *Rhipicephalus (Boophilus) microplus*. [Clin Vaccine Immunol. 16: 1074-86.](#)
11. Woolums, A.R. *et al.* (2013) Effect of calf age and administration route of initial multivalent modified-live virus vaccine on humoral and cell-mediated immune responses following subsequent administration of a booster vaccination at weaning in beef calves. [Am J Vet Res. 74: 343-54.](#)
12. McInnes, E. *et al.* (1999) Phenotypic analysis of local cellular responses in calves infected with bovine respiratory syncytial virus. [Immunology. 96 \(3\): 396-403.](#)
13. Maślanka, T. *et al.* (2012) The presence of CD25 on bovine WC1+ $\gamma\delta$ T cells is positively correlated with their production of IL-10 and TGF- β , but not IFN- γ . [Pol J Vet Sci. 15: 11-20.](#)
14. Menge, C. *et al.* (2003) Verotoxin 1 from *Escherichia coli* affects Gb3/CD77+ bovine lymphocytes independent of interleukin-2, tumor necrosis factor-alpha, and interferon-alpha. [Exp Biol Med \(Maywood\). 228: 377-86.](#)
15. Menge, C. *et al.* (1999) Shiga toxin 1 from *Escherichia coli* blocks activation and proliferation of bovine lymphocyte subpopulations *in vitro*. [Infect Immun. 67: 2209-17.](#)
16. Constantinoiu, C.C. *et al.* (2010) Local immune response against larvae of *Rhipicephalus (Boophilus) microplus* in *Bos taurus indicus* and *Bos taurus taurus* cattle. [Int J Parasitol. 40: 865-75.](#)
17. Maślanska, T. and Jaroszewski, J.J. (2013) In vitro effects of meloxicam on the number, Foxp3 expression, production of selected cytokines, and apoptosis of bovine CD25+CD4+ and CD25-CD4+ cells. [J Vet Sci. 14 \(2\): 125-34.](#)
18. Brodzki, P. *et al.* (2014) Phenotyping of leukocytes and granulocyte and monocyte phagocytic activity in the peripheral blood and uterus of cows with endometritis. [Theriogenology. 82 \(3\): 403-10.](#)
19. Zoldan, K. *et al.* (2014) Increase of CD25 expression on bovine neutrophils correlates with disease severity in post-partum and early lactating dairy cows. [Dev Comp Immunol. 47 \(2\): 254-63.](#)
20. Kang, H.J. *et al.* (2016) Effects of Ambient Temperature on Growth Performance, Blood Metabolites, and Immune Cell Populations in Korean Cattle Steers. [Asian-Australas J Anim Sci. 29 \(3\): 436-43.](#)
21. Schmidt, N. *et al.* (2018) Decreased STEC shedding by cattle following passive and active vaccination based on recombinant *Escherichia coli* Shiga toxoids. [Vet Res. 49 \(1\): 28.](#)
22. Higgins, J.L. *et al.* (2018) Cell mediated immune response in goats after experimental challenge with the virulent *Brucella melitensis* strain 16M and the reduced virulence strain Rev. 1. [Vet Immunol Immunopathol. 202: 74-84.](#)
23. Brodzki, P. *et al.* (2019) Selected leukocyte subpopulations in peripheral blood and uterine washings in cows before and after intrauterine administration of cefapirin and methisoprinol. [Anim Sci J. Nov 06 \[Epub ahead of print\].](#)

Storage

Store at +4°C or at -20°C if preferred.
Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2430GA 10040
--------------------------------------	---

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M366894:200529'

Printed on 14 Oct 2024