

Datasheet: MCA2433F BATCH NUMBER 158799

Description:	MOUSE ANTI BOVINE CD44:FITC
Specificity:	CD44
Other names:	H-CAM
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	IL-A118
lsotype:	lgG1
Quantity:	0.1 mg

Product Details

Applications	pplications This product has been reported to work in the following applications. This information					
	derived from testing within our laboratories, peer-reviewed publications or personal					
	communications from the originators. Please refer to references indicated for further					
	information. For general protocol recommendations, please visit <u>www.bio-</u> rad-antibodies.com/protocols.					
	· · · · · ·	Yes No	Not Determined	Suggested Dilution		
	Flow Cytometry	•		Neat - 1/10		
	Where this antibody has not been tested for use in a particular technique this does not					
	necessarily exclude its	s use in such procedu	ures. Suggested work	king dilutions are given as		
	a guide only. It is reco	mmended that the us	er titrates the antiboo	dy for use in their own		
	system using appropri	ate negative/positive	controls.			
	, , , , , , , , , , , , , , , , , , , ,	U				
Target Species	Bovine					
Species Cross	Reacts with: Camel					
Reactivity	N.B. Antibody reactivity and working conditions may vary between species. Cros					
	reactivity is derived fro	om testing within our	aboratories, peer-rev	viewed publications or		
	personal communications from the originators. Please refer to references in					
	further information.					
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid					
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)	_		
	FITC	490	525	_		
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture					

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
External Database Links	UniProt: Q29423 Related reagents Entrez Gene: 281057 CD44 Related reagents
RRID	AB_1604795
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the X63.Ag8.853 myeloma cell line.
Specificity	Mouse anti Bovine CD44 antibody, clone IL-A118 recognizes bocine Phagocytic Glycoprotein-1 (PGP-1), also known as CD44, Hermes antigen, Extracellular matrix receptor III or HUTCH-1. Bovine CD44 is a 346 amino acid ~90 kDa type I single pass transmembrane glycoprotein containing a single Link domain, responsible for hyaluronan binding. Bovine CD44 is expressed by a wide range of bovine cells, including peripheral T and B lymphocytes, monocytes, granulocytes, platelets and early erythroid cells.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^{6} cells in 100ul
References	 Naessens, J. & Nthale, J. (1993) Biochemical characterization of three non-lineage antigens defined by workshop antibodies. <u>Vet Immunol Immunopathol. 39 (1-3): 217-23.</u> Naessens, J. <i>et al.</i> (1993) Cross-reactivity of workshop antibodies with cells from domestic and wild ruminants. <u>Vet Immunol Immunopathol. 39 (1-3): 283-90.</u> Howard, C.J. & Naessens, J. (1993) Summary of workshop findings for cattle (tables 1 and 2). <u>Vet Immunol Immunopathol. 39 (1-3): 25-47.</u> Menge C <i>et al.</i> (2004) Bovine ileal intraepithelial lymphocytes represent target cells for Shiga toxin 1 from <i>Escherichia coli.</i> <u>Infect Immun. 72 (4): 1896-905.</u> de Moraes, C.N. <i>et al.</i> (2016) Bovine endometrial cells: a source of mesenchymal stem/progenitor cells. <u>Cell Biol Int. 40 (12): 1332-9.</u> de Moraes, C.N. <i>et al.</i> (2017) Shotgun proteomic analysis of the secretome of bovine endometrial mesenchymal progenitor/stem cells challenged or not with bacterial lipopolysaccharide. <u>Vet Immunol Immunopathol. 187: 42-7.</u> Lee, J. <i>et al.</i> (2020) Bovine tongue epithelium-derived cells: A new source of bovine mesenchymal stem cells. <u>Biosci Rep. 40 (4): BSR20181829.</u> Molinos, M. <i>et al.</i> (2023) Alterations of bovine nucleus pulposus cells with aging. <u>Aging Cell. 22 (8): e13873.</u> Ferreira, J.R. <i>et al.</i> (2024) Dynamics of CD44(+) bovine nucleus pulposus cells with inflammation. <u>Sci Rep. 14 (1): 9156.</u>

	10. Barcelos, S.M. <i>et al.</i> (2024) Extracellular vesicles derived fi mesenchymal stromal cells enhance <i>in vitro</i> embryo production <u>Cytotherapy. 26 (10): 1141-51.</u>	rom bovine adipose-derived n from lesioned ovaries.	
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.		
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.		
Guarantee	12 months from date of despatch		
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA2433F 10041		
Regulatory	For research purposes only		

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL: FITC (MCA928F)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad	.com	Email: antibody_sales_uk@bio-rad	.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M385690:210513'

Printed on 08 Nov 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint