

## Datasheet: MCA2433F

|                      |                             |
|----------------------|-----------------------------|
| <b>Description:</b>  | MOUSE ANTI BOVINE CD44:FITC |
| <b>Specificity:</b>  | CD44                        |
| <b>Other names:</b>  | H-CAM                       |
| <b>Format:</b>       | FITC                        |
| <b>Product Type:</b> | Monoclonal Antibody         |
| <b>Clone:</b>        | IL-A118                     |
| <b>Isotype:</b>      | IgG1                        |
| <b>Quantity:</b>     | 0.1 mg                      |

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

|                | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪   |    |                | Neat - 1/10        |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

### Target Species

Bovine

### Species Cross Reactivity

Reacts with: Camel

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

### Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

### Max Ex/Em

| Fluorophore | Excitation Max (nm) | Emission Max (nm) |
|-------------|---------------------|-------------------|
| FITC        | 490                 | 525               |

### Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

### Buffer Solution

Phosphate buffered saline

|                                       |   |
|---------------------------------------|---|
| <b>Preservative Stabilisers</b>       | 0.09% Sodium Azide<br>1% Bovine Serum Albumin   |
| <b>Approx. Protein Concentrations</b> | IgG concentration 0.1 mg/ml   |
| <b>External Database Links</b>        | <p><b>UniProt:</b><br/> <a href="#">Q29423</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b><br/> <a href="#">281057</a> CD44    <a href="#">Related reagents</a></p>  |
| <b>RRID</b>                           | AB_1604795  |
| <b>Fusion Partners</b>                | Spleen cells from immunised BALB/c mice were fused with cells of the X63.Ag8.853 myeloma cell line.   |
| <b>Specificity</b>                    | <b>Mouse anti Bovine CD44 antibody, clone IL-A118</b> recognizes bovine Phagocytic Glycoprotein-1 (PGP-1), also known as CD44, Hermes antigen, Extracellular matrix receptor III or HUTCH-1. Bovine CD44 is a 346 amino acid ~90 kDa type I single pass transmembrane glycoprotein containing a single <a href="#">Link domain</a> , responsible for hyaluronan binding. Bovine CD44 is expressed by a wide range of bovine cells, including peripheral T and B lymphocytes, monocytes, granulocytes, platelets and early erythroid cells.  |
| <b>Flow Cytometry</b>                 | Use 10ul of the suggested working dilution to label $1 \times 10^6$ cells in 100ul  |
| <b>References</b>                     | <ol style="list-style-type: none"> <li>Naessens, J. &amp; Nthale, J. (1993) Biochemical characterization of three non-lineage antigens defined by workshop antibodies. <a href="#">Vet Immunol Immunopathol. 39 (1-3): 217-23.</a></li> <li>Menge C <i>et al.</i> (2004) Bovine ileal intraepithelial lymphocytes represent target cells for Shiga toxin 1 from <i>Escherichia coli</i>. <a href="#">Infect Immun. 72 (4): 1896-905.</a></li> <li>Naessens, J. <i>et al.</i> (1993) Cross-reactivity of workshop antibodies with cells from domestic and wild ruminants. <a href="#">Vet Immunol Immunopathol. 39 (1-3): 283-90.</a></li> <li>Howard, C.J. &amp; Naessens, J. (1993) Summary of workshop findings for cattle (tables 1 and 2). <a href="#">Vet Immunol Immunopathol. 39 (1-3): 25-47.</a></li> <li>de Moraes, C.N. <i>et al.</i> (2016) Bovine endometrial cells: a source of mesenchymal stem/progenitor cells. <a href="#">Cell Biol Int. 40 (12): 1332-9.</a></li> <li>de Moraes, C.N. <i>et al.</i> (2017) Shotgun proteomic analysis of the secretome of bovine endometrial mesenchymal progenitor/stem cells challenged or not with bacterial lipopolysaccharide. <a href="#">Vet Immunol Immunopathol. Mar 27 [Epub ahead of print]</a></li> <li>Lee, J. <i>et al.</i> (2020) Bovine tongue epithelium-derived cells: A new source of bovine mesenchymal stem cells. <a href="#">Biosci Rep. 40 (4): BSR20181829.</a></li> </ol> |
| <b>Storage</b>                        | <p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in</p>   |

frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

---

|                  |                                 |
|------------------|---------------------------------|
| <b>Guarantee</b> | 12 months from date of despatch |
|------------------|---------------------------------|

---

|                                      |  |
|--------------------------------------|--|
| <b>Health And Safety Information</b> | Material Safety Datasheet documentation #10041 available at:<br><a href="https://www.bio-rad-antibodies.com/SDS/MCA2433F">https://www.bio-rad-antibodies.com/SDS/MCA2433F</a><br>10041 |
|--------------------------------------|--|

---

|                   |                            |
|-------------------|----------------------------|
| <b>Regulatory</b> | For research purposes only |
|-------------------|----------------------------|

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M385690:210513'

**Printed on 25 Mar 2023**

---

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)