

Datasheet: MCA2434GA

Description:	MOUSE ANTI BOVINE CD45RO
Specificity:	CD45RO
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	IL-A116
Isotype:	IgG3
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/50
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Bovine
Species Cross Reactivity	Reacts with: Goat N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml

Immunogen	Bovine peripheral blood mononuclear cells.
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the X63.Ag8.653 myeloma cell line.
Specificity	<p>Mouse anti Bovine CD45RO, clone IL-A116 recognises the bovine homologue of the human CD45RO cell surface antigen.</p> <p>CD45, also known as Leucocyte Common Antigen or LCA, occurs in a number of isoforms, clone IL-A116 is specific for the low molecular weight isoform termed CD45RO, the isoform associated with expression on memory T-cells. Bovine CD45RO is expressed by monocytes, granulocytes and subsets of thymocytes, CD4+ T cells and CD8+ T cells. Studies utilizing clone IL-A116 have demonstrated that the percentage of CD45RO⁺ CD8⁺ T cells increase from approximately 5% in neonatal calves to approximately 35% in adult cattle over the age of 5 years (Hogg et al. 2011). It has been demonstrated that mouse anti bovine CD45RO, clone IL-A116 immunoprecipitates a molecule of ~180 kDa (Bembridge et al. 1995) which is analogous to the molecular weight of human and mouse CD45RO.</p> <p>Mouse anti Bovine CD45RO, clone IL-A116 has been demonstrated to recognise the CD45RO cell surface antigen by flow cytometry in both European cattle, <i>Bos taurus</i>, and in Zebu, <i>B.indicus</i> (Bembridge et al. 1995).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul
References	<ol style="list-style-type: none"> 1. Bembridge, G.P. <i>et al.</i> (1995) CD45RO expression on bovine T cells: relation to biological function. Immunology. 86 (4): 537-44. 2. Hogg, A.E. <i>et al.</i> (2011) Characterization of age-related changes in bovine CD8+ T-cells. Vet Immunol Immunopathol. 140 (1-2): 47-54. 3. Whelan, A.O. <i>et al.</i> (2011) Development of an antibody to bovine IL-2 reveals multifunctional CD4 T(EM) cells in cattle naturally infected with bovine tuberculosis. PLoS One. 6 (12): e29194. 4. Howard, C.J. & Naessens, J. (1993) Summary of workshop findings for cattle (tables 1 and 2). Vet Immunol Immunopathol. 39 (1-3): 25-47. 5. Sopp, P. & Howard, C.J. (2001) IFN gamma and IL-4 production by CD4, CD8 and WC1 gamma delta TCR(+) T cells from cattle lymph nodes and blood. Vet Immunol Immunopathol. 81 (1-2): 85-96. 6. McInnes, E. <i>et al.</i> (1999) Phenotypic analysis of local cellular responses in calves infected with bovine respiratory syncytial virus. Immunology. 96 (3): 396-403. 7. Bembridge, G.P. <i>et al.</i> (1993) Comparison of monoclonal antibodies with potential specificity for restricted isoforms of the leukocyte common antigen (CD45R). Vet Immunol Immunopathol. 39 (1-3): 129-36. 8. Naessens, J. <i>et al.</i> (1993) Cross-reactivity of workshop antibodies with cells from domestic and wild ruminants. Vet Immunol Immunopathol. 39 (1-3): 283-90. 9. Schmidt, N. <i>et al.</i> (2018) Decreased STEC shedding by cattle following passive and active vaccination based on recombinant <i>Escherichia coli</i> Shiga toxoids. Vet Res. 49 (1): 28.
Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>Storage in frost-free freezers is not recommended.</p> <p>This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
Guarantee	18 months from date of despatch

**Health And Safety
Information**

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Goat Anti Mouse IgG (STAR77...) [HRP](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Rabbit Anti Mouse IgG (STAR8...) [DyLight®800](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Human Anti Mouse IgG3 (HCA039...) [FITC](#), [HRP](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®680](#),
[DyLight®800](#), [FITC](#), [HRP](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M334627:181203'

Printed on 20 May 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)