

# Datasheet: MCA2434PE

Description:	MOUSE ANTI BOVINE CD45RO:RPE
Specificity:	CD45RO
Format:	RPE
<b>Product Type:</b>	Monoclonal Antibody
Clone:	IL-A116
Isotype:	lgG3
Quantity:	100 TESTS

#### **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Bovine			
Species Cross	Reacts with: Goat			
Reactivity	reactivity is derived	from testing within our I	ons may vary between species. Cros aboratories, peer-reviewed publicatio ors. Please refer to references indicat	ns or
Product Form	Purified IgG conjuga	ated to R. Phycoerythrin	(RPE) - lyophilized	
Reconstitution	Reconstitute with 1.	0 ml distilled water		
	Care should be take	en during reconstitution	as the protein may appear as a film a	at the
	bottom of the vial. B	Bio-Rad recommend that	t the vial is gently mixed after reconst	titution
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)	
	RPE 488nm laser	496	578	

# supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> ) 1% Bovine Serum Albumin 5% Sucrose
Immunogen	Bovine peripheral blood monouclear cells.
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the X63.Ag8.653 myeloma cell line.
Specificity	Mouse anti Bovine CD45RO, clone IL-A116 recognizes the bovine homologue of the human CD45RO cell surface antigen.
	CD45, also known as Leucocyte Common Antigen or LCA, occurs in a number of isoforms, clone IL-A116 is specific for the low molecular weight isoform termed CD45RO, the isoform associated with expression on memory T-cells. Bovine CD45RO is expressed by monocytes, granulocytes and subsets of thymocytes, CD4+ T cells and CD8+ T cells. CD45RO+ CD8+ T cells increase from approximately 5% in neonatal calves to approximately 35% in adult cattle over the age of 5 years (Hogg <i>et al.</i> 2011). Mouse anti Bovine CD45RO, clone IL-A116 immunoprecipitates a molecule of ~180 kDa (Bembridge <i>et al.</i> 1995), analogus to the molecular weight of human and mouse CD45RO.  Mouse anti Bovine CD45RO, clone IL-A116 recognizes the CD45RO cell surface antigen by flow cytometry in both European cattle, <i>Bos taurus</i> , and in Zebu, <i>B.indicus</i> (Bembridge <i>et al.</i> 1995).
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul
References	1. Howard, C.J. & Naessens, J. (1993) Summary of workshop findings for cattle (tables 1 and 2). Vet Immunol Immunopathol. 39 (1-3): 25-47.  2. Naessens, J. et al. (1993) Cross-reactivity of workshop antibodies with cells from domestic and wild ruminants. Vet Immunol Immunopathol. 39 (1-3): 283-90.  3. Bembridge, G.P. et al. (1993) Comparison of monoclonal antibodies with potential specificity for restricted isoforms of the leukocyte common antigen (CD45R). Vet Immunol Immunopathol. 39 (1-3): 129-36.  4. Bembridge, G.P. et al. (1995) CD45RO expression on bovine T cells: relation to biological function. Immunology. 86 (4): 537-44.  5. Mcinnes, E. et al. (1999) Phenotypic analysis of local cellular responses in calves infected with bovine respiratory syncytial virus. Immunology. 96 (3): 396-403.  6. Sopp, P. & Howard, C.J. (2001) IFN gamma and IL-4 production by CD4, CD8 and WC1 gamma delta TCR(+) T cells from cattle lymph nodes and blood. Vet Immunol Immunopathol. 81 (1-2): 85-96.

8. Whelan, A.O. et al. (2011) Development of an antibody to bovine IL-2 reveals

multifunctional CD4 T(EM) cells in cattle naturally infected with bovine tuberculosis. PLoS One. 6 (12): e29194.

- 9. Schmidt, N. et al. (2018) Decreased STEC shedding by cattle following passive and active vaccination based on recombinant Escherichia coli Shiga toxoids. Vet Res. 49 (1): 28.
- 10. Arrieta-Villegas, C. et al. (2020) Immunogenicity and Protection against Mycobacterium caprae Challenge in Goats Vaccinated with BCG and Revaccinated after One Year. Vaccines (Basel). 8 (4): 751.
- 11. Wherry, T.L.T. et al. (2022) Effects of 1,25-Dihydroxyvitamin D<sub>3</sub> and 25-Hydroxyvitamin D<sub>3</sub> on PBMCs From Dairy Cattle Naturally Infected With Mycobacterium avium subsp. paratuberculosis. Front Vet Sci. 9: 830144.
- 12. Hidalgo-Ruiz, M. et al. (2022) Babesia bovis AMA-1, MSA-2c and RAP-1 contain conserved B and T-cell epitopes, which generate neutralizing antibodies and a long-lasting Th1 immune response in vaccinated cattle. Vaccine. S0264-410X(22)00049-4.
- 13. Seemann, L. et al. (2024) Dietary L-carnitine supplementation modifies blood parameters of mid-lactating dairy cows during standardized lipopolysaccharide-induced inflammation. Front Immunol. 15: 1390137.

#### Storage

This product is shipped at ambient temperature.

Prior to reconstitution store at +4°C.

Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2434PE">https://www.bio-rad-antibodies.com/SDS/MCA2434PE</a> 20487
Regulatory	For research purposes only

America

North & South Tel: +1 800 265 7376 Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_us@bio-rad.com

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M440627:250523'

### Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint