

## Datasheet: MCA2434PE

<b>Description:</b>	MOUSE ANTI BOVINE CD45RO:RPE
<b>Specificity:</b>	CD45RO
<b>Format:</b>	RPE
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	IL-A116
<b>Isotype:</b>	IgG3
<b>Quantity:</b>	100 TESTS

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

### Target Species

Bovine

### Species Cross Reactivity

Reacts with: Goat

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

### Product Form

Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized

### Reconstitution

Reconstitute with 1.0 ml distilled water

Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.

### Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
RPE 488nm laser	496	578

### Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture

supernatant

---

**Buffer Solution** Phosphate buffered saline

---

**Preservative** 0.09% Sodium Azide (NaN<sub>3</sub>)  
**Stabilisers** 1% Bovine Serum Albumin  
5% Sucrose

---

**Immunogen** Bovine peripheral blood mononuclear cells.

---

**Fusion Partners** Spleen cells from immunised BALB/c mice were fused with cells of the X63.Ag8.653 myeloma cell line.

---

**Specificity** **Mouse anti Bovine CD45RO, clone IL-A116** recognises the bovine homologue of the human CD45RO cell surface antigen.

CD45, also known as Leucocyte Common Antigen or LCA, occurs in a number of isoforms, clone IL-A116 is specific for the low molecular weight isoform termed CD45RO, the isoform associated with expression on memory T-cells. Bovine CD45RO is expressed by monocytes, granulocytes and subsets of thymocytes, CD4<sup>+</sup> T cells and CD8<sup>+</sup> T cells. Studies utilizing clone IL-A116 have demonstrated that the percentage of CD45RO<sup>+</sup> CD8<sup>+</sup> T cells increase from approximately 5% in neonatal calves to approximately 35% in adult cattle over the age of 5 years ([Hogg et al. 2011](#)). It has been demonstrated that mouse anti bovine CD45RO, clone IL-A116 immunoprecipitates a molecule of ~180 kDa ([Bembridge et al. 1995](#)) which is analogous to the molecular weight of human and mouse CD45RO.

Mouse anti Bovine CD45RO, clone IL-A116 has been demonstrated to recognise the CD45RO cell surface antigen by flow cytometry in both European cattle, *Bos taurus*, and in Zebu, *B.indicus* ([Bembridge et al. 1995](#)).

---

**Flow Cytometry** Use 10ul of the suggested working dilution to label 1x10<sup>6</sup> cells in 100ul

---

- References**
1. Bembridge, G.P. *et al.* (1995) CD45RO expression on bovine T cells: relation to biological function. [Immunology. 86 \(4\): 537-44.](#)
  2. Hogg, A.E. *et al.* (2011) Characterization of age-related changes in bovine CD8<sup>+</sup> T-cells. [Vet Immunol Immunopathol. 140 \(1-2\): 47-54.](#)
  3. Whelan, A.O. *et al.* (2011) Development of an antibody to bovine IL-2 reveals multifunctional CD4 T(EM) cells in cattle naturally infected with bovine tuberculosis. [PLoS One. 6 \(12\): e29194.](#)
  4. Howard, C.J. & Naessens, J. (1993) Summary of workshop findings for cattle (tables 1 and 2). [Vet Immunol Immunopathol. 39 \(1-3\): 25-47.](#)
  5. Sopp, P. & Howard, C.J. (2001) IFN gamma and IL-4 production by CD4, CD8 and WC1 gamma delta TCR(+) T cells from cattle lymph nodes and blood. [Vet Immunol Immunopathol. 81 \(1-2\): 85-96.](#)
  6. McInnes, E. *et al.* (1999) Phenotypic analysis of local cellular responses in calves infected with bovine respiratory syncytial virus. [Immunology. 96 \(3\): 396-403.](#)
  7. Bembridge, G.P. *et al.* (1993) Comparison of monoclonal antibodies with potential

specificity for restricted isoforms of the leukocyte common antigen (CD45R). [Vet Immunol Immunopathol. 39 \(1-3\): 129-36.](#)

8. Naessens, J. *et al.* (1993) Cross-reactivity of workshop antibodies with cells from domestic and wild ruminants. [Vet Immunol Immunopathol. 39 \(1-3\): 283-90.](#)

9. Schmidt, N. *et al.* (2018) Decreased STEC shedding by cattle following passive and active vaccination based on recombinant *Escherichia coli* Shiga toxoids. [Vet Res. 49 \(1\): 28.](#)

10. Arrieta-Villegas, C. *et al.* (2020) Immunogenicity and Protection against *Mycobacterium caprae* Challenge in Goats Vaccinated with BCG and Revaccinated after One Year. [Vaccines \(Basel\). 8 \(4\): 751.](#)

---

**Storage**

Prior to reconstitution store at +4°C.

Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety Information**

Material Safety Datasheet documentation #20487 available at:  
20487: <https://www.bio-rad-antibodies.com/uploads/MSDS/20487.pdf>

---

**Regulatory**

For research purposes only

---

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M375492:210104'

**Printed on 30 Mar 2021**

---

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)