# Datasheet: MCA2436GA BATCH NUMBER 1114

Description:	MOUSE ANTI BOVINE CD80
Specificity:	CD80
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	IL-A159
lsotype:	lgG1
Quantity:	0.1 mg

# **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .				
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry	•			1/25 - 1/200
	Immunohistology - Frozen				
	Immunohistology - Paraffin				
	ELISA				
	Immunoprecipitation			•	
	Western Blotting				
Target Species	necessarily exclude its us a guide only. It is recomn system using appropriate Bovine	nended that	at the use	r titrates the antibody	•
Species Cross Reactivity	Reacts with: Sheep <b>N.B.</b> Antibody reactivity a reactivity is derived from personal communications further information.	testing wit	hin our la	boratories, peer-revie	wed publications or
Product Form	Purified IgG - liquid				
Preparation	Purified IgG prepared by supernatant	affinity ch	romatogra	aphy on Protein A fror	n tissue culture

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> )
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
External Database Links	UniProt: 046405 Related reagents
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the X63.Ag8.653 myeloma cell line.
Specificity	<b>Mouse anti Bovine CD80 antibody, clone IL-A159</b> recognizes the bovine CD80 cell surface antigen, expressed by dendritic cells, activated macrophages and activated B cells. CD80 plays a key role in co-stimulation of T cells during the primary immune response.
Flow Cytometry	Use 10ul of the suggested working dilution to label $1 \times 10^6$ cells in 100ul.
References	<ol> <li>Langelaar, M.F. <i>et al.</i> (2005) <i>Mycobacterium avium</i> ssp. <i>paratuberculosis</i> recombinant heat shock protein 70 interaction with different bovine antigen-presenting cells. <u>Scand J</u> <u>Immunol. 61 (3): 242-50.</u></li> <li>Rhodes, S.G. <i>et al.</i> (2003) 1,25-dihydroxyvitamin D3 and development of tuberculosis in cattle. <u>Clin Diagn Lab Immunol. 10 (6): 1129-35.</u></li> <li>Glew, E.J. <i>et al.</i> (2003) Differential effects of bovine viral diarrhoea virus on monocytes and dendritic cells. <u>J Gen Virol. 84 (Pt 7): 1771-80.</u></li> <li>Epardaud, M. <i>et al.</i> (2004) Enrichment for a CD26hi SIRP- subset in lymph dendritic cells from the upper aero-digestive tract. <u>J Leukoc Biol. 76 (3): 553-61.</u></li> <li>Bonneau, M. <i>et al.</i> (2006) Migratory monocytes and granulocytes are major lymphatic carriers of Salmonella from tissue to draining lymph node. <u>J Leukoc Biol. 79: 268-76.</u></li> <li>Hart, J. <i>et al.</i> (2011) <i>Theileria annulata</i>-transformed cell lines are efficient antigen- presenting cells for <i>in vitro</i> analysis of CD8 T cell responses to bovine herpesvirus-1. <u>Vet Res. 42: 119.</u></li> <li>Ikebuchi, R. <i>et al.</i> (2013) Blockade of bovine PD-1 increases T cell function and inhibits bovine leukemia virus expression in B cells <i>in vitro</i>. <u>Vet Res. 44: 59.</u></li> <li>Totté P <i>et al.</i> (2015) Free exopolysaccharide from <i>Mycoplasma mycoides</i> subsp. <i>mycoides</i> possesses anti-inflammatory properties. <u>Vet Res. 46 (1): 122.</u></li> <li>Corripio-miyar, Y. <i>et al.</i> (2017) 1,25-Dihydroxyvitamin D3 modulates the phenotype and function of Monocyte derived dendritic cells in cattle <u>BMC Veterinary Research. 13 (1)</u> [Epub ahead of print].</li> </ol>
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may

	denature the antibody. Should this product contain a precipitate we reminicrocentrifugation before use.	ecommend
Guarantee	12 months from date of despatch	
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2436GA">https://www.bio-rad-antibodies.com/SDS/MCA2436GA</a> 10040	
Regulatory	For research purposes only	

## Related Products

## **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12)	RPE		
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>			
Goat Anti Mouse IgG (STAR76)	RPE		
Rabbit Anti Mouse IgG (STAR13)	HRP		
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>		
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,		
	DyLight®650, DyLight®680, DyLight®800,		
	<u>FITC, HRP</u>		
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>		
Goat Anti Mouse IgG (STAR77)	HRP		
Goat Anti Mouse IgG (Fc) (STAR120)	<u>FITC, HRP</u>		

#### MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad	.com	Email: antibody_sales_uk@bio-rac	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M366903:200529'

#### Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint