

Datasheet: MCA2437A647

### **BATCH NUMBER 163362**

Description:	MOUSE ANTI BOVINE CD86:Alexa Fluor® 647		
Specificity:	CD86		
Format:	ALEXA FLUOR® 647		
Product Type:	Monoclonal Antibody		
Clone:	IL-A190		
Isotype:	IgG1		
Quantity:	100 TESTS/1ml		

## **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Bovine			
Species Cross	Reacts with: Sheep	)		
Reactivity	reactivity is derived	tivity and working condit I from testing within our I cations from the originato	aboratories, peer-re	viewed publications o
Product Form	Purified IgG conjugated to Alexa Fluor 647 - liquid			
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm	)
	Alexa Fluor®647	650	665	
Preparation	Purified IgG prepar supernatant	ed by affinity chromatog	raphy on Protein A	from tissue culture
Buffer Solution	Phosphate buffered	d 1:		

Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> ) 1% Bovine Serum Albumin			
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml			
External Database Links	UniProt:  Q1JPC5 Related reagents			
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the X63.Ag8.653 myeloma cell line.			
Specificity	Mouse anti Bovine CD86 antibody, cllone IL-A190 recognizes the bovine CD86 cell surface antigen, expressed by dendritic cells, activated macrophages and activated B cells. CD86 plays an important role in co-stimulation of T cells in the primary immune response.			
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul			
References	<ol> <li>Norimatsu, M. <i>et al.</i> (2003) Differential response of bovine monocyte-derived macrophages and dendritic cells to infection with Salmonella typhimurium in a low-dose model in vitro. <u>Immunology. 108: 55-61.</u></li> <li>Glew, E.J. <i>et al.</i> (2003) Differential effects of bovine viral diarrhoea virus on monocytes and dendritic cells. <u>J Gen Virol. 84 (Pt 7): 1771-80.</u></li> <li>Rhodes, S.G. <i>et al.</i> (2003) 1,25-dihydroxyvitamin D3 and development of tuberculosis in cattle. <u>Clin Diagn Lab Immunol. 10 (6): 1129-35.</u></li> <li>Epardaud, M. <i>et al.</i> (2004) Enrichment for a CD26hi SIRP- subset in lymph dendritic cells from the upper aero-digestive tract. <u>J Leukoc Biol. 76 (3): 553-61.</u></li> <li>Langelaar, M.F. <i>et al.</i> (2005) <i>Mycobacterium avium ssp. paratuberculosis</i> recombinant</li> </ol>			

Immunol. 61: 242-50
6. Bonneau, M. *et al.* (2006) Migratory monocytes and granulocytes are major lymphatic carriers of Salmonella from tissue to draining lymph node. <u>J Leukoc Biol. 79: 268-76.</u>

heat shock protein 70 interaction with different bovine antigen-presenting cells. Scand J

- 7. Pascale, F. *et al.* (2008) Plasmacytoid dendritic cells migrate in afferent skin lymph. <u>J</u> Immunol. 180: 5963-72.
- 8. Hemati, B. *et al.* (2009) Bluetongue virus targets conventional dendritic cells in skin lymph. J Virol. 83: 8789-99.
- 9. Ruscanu, S. *et al.* (2012) The double-stranded RNA bluetongue virus induces type I interferon in plasmacytoid dendritic cells via a MYD88-dependent TLR7/8-independent signaling pathway. J Virol. 2012 May;86: 5817-28.
- 10. Mauro, A. *et al.* (2016) M1 and M2 macrophage recruitment during tendon regeneration induced by amniotic epithelial cell allotransplantation in ovine. Res Vet Sci. 105: 92-102.
- 11. Corripio-miyar, Y. *et al.* (2017) 1,25-Dihydroxyvitamin D3 modulates the phenotype and function of Monocyte derived dendritic cells in cattle. <u>BMC Vet Res. 13 (1): 390.</u>
- 12. Liu, J. et al. (2020) *Theileria annulata*. transformation altered cell surface molecules expression and endocytic function of monocyte-derived dendritic cells. <u>Ticks Tick Borne</u>

#### Dis. 11 (3): 101365.

- 13. Marzo, S. *et al.* (2021) Characterisation of dendritic cell frequency and phenotype in bovine afferent lymph reveals kinetic changes in costimulatory molecule expression <u>Vet Immunol Immunopathol.</u> 19 Nov: 110363.
- 14. Stabel, J.R. *et al.* (2022) B cell phenotypes and maturation states in cows naturally infected with *Mycobacterium avium* subsp. *Paratuberculosis*. <u>PLoS One. 17 (12):</u> e0278313.
- 15. Russo, V. *et al.* (2022) Tendon Healing Response Is Dependent on Epithelial-Mesenchymal-Tendon Transition State of Amniotic Epithelial Stem Cells. <u>Biomedicines. 10</u> (5): 1177.
- 16. Wherry, T.L.T. *et al.* (2022) Effects of 1,25-Dihydroxyvitamin D<sub>3</sub> and 25-Hydroxyvitamin D<sub>3</sub> on PBMCs From Dairy Cattle Naturally Infected With *Mycobacterium avium* subsp. *paratuberculosis*. Front Vet Sci. 9: 830144.
- 17. Zenobi, M.G. *et al.* (2020) Effect of prepartum energy intake and supplementation with ruminally protected choline on innate and adaptive immunity of multiparous Holstein cows. <u>J Dairy Sci. 103 (3): 2200-16.</u>
- 18. Zhou, X. *et al.* (2022) Dusp6 deficiency attenuates neutrophil-mediated cardiac damage in the acute inflammatory phase of myocardial infarction. <u>Nat Commun. 13 (1):</u> 6672.
- 19. Kornuta, C.A. *et al.* (2025) Galectin-8 and GEL01 as potential adjuvants to enhance the immune response induced by a DNA vaccine against bovine alphaherpesvirus Type-1. <u>Virology. 604: 110402.</u>

### Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

### Guarantee

12 months from date of despatch

### Acknowledgements

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchased product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

# Health And Safety Information

Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2437A647">https://www.bio-rad-antibodies.com/SDS/MCA2437A647</a> 10041

### Regulatory

For research purposes only

# **Related Products**

# **Recommended Negative Controls**

### MOUSE IgG1 NEGATIVE CONTROL: Alexa Fluor® 647 (MCA928A647)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M387825:210726'

### Printed on 24 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint