

Datasheet: MCA2440P

Description:	MOUSE ANTI BOVINE IgG1:HRP
Specificity:	IgG1
Format:	HRP
Product Type:	Monoclonal Antibody
Clone:	IL-A60
Isotype:	IgG1
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			1/100 - 1/1000
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Bovine
Product Form	Purified IgG conjugated to Horseradish Peroxidase (HRP) - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.01% Thiomersal
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml

Immunogen	Purified bovine immunoglobulin.
RRID	AB_1605095
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the X63.Ag.653 myeloma cell line.
Specificity	Mouse anti Bovine IgG1 antibody, clone IL-A60 recognizes bovine IgG1. Mouse anti Bovine IgG1 antibody, clone IL-A60 immunoprecipitates a protein band of 55-59 kDa, consistent with the heavy chain of bovine IgG1 (Campbell et al. 1998).
References	<ol style="list-style-type: none"> Williams, D.J. <i>et al.</i> (1996) The role of anti-variable surface glycoprotein antibody responses in bovine trypanotolerance. Parasite Immunol. 18 (4): 209-18. Campbell, J.D. <i>et al.</i> (1998) A novel cell surface proliferation-associated marker expressed on T cells and up-regulated on germinal center B cells. J Leukoc Biol. 63 (5): 567-74. Hecker YP <i>et al.</i> (2014) A <i>Neospora caninum</i> vaccine using recombinant proteins fails to prevent foetal infection in pregnant cattle after experimental intravenous challenge. Vet Immunol Immunopathol. 162 (3-4): 142-53. Dorneles, E.M. <i>et al.</i> (2015) Immune Response of Calves Vaccinated with <i>Brucella abortus</i> S19 or RB51 and Revaccinated with RB51. PLoS One. 10 (9): e0136696. Jaramillo, J.O. <i>et al.</i> (2019) Immunisation of cattle against <i>Babesia bovis</i>. combining a multi-epitope modified vaccinia Ankara virus and a recombinant protein induce strong Th1 cell responses but fails to trigger neutralising antibodies required for protection. Ticks Tick Borne Dis. 10 (6): 101270. Pereyra, R. <i>et al.</i> (2019) Evidence of reduced vertical transmission of <i>Neospora caninum</i>. associated with higher IgG1 than IgG2 serum levels and presence of IFN-γ in non-aborting chronically infected cattle under natural condition. Vet Immunol Immunopathol. 208: 53-57. Hecker, Y.P. <i>et al.</i> (2019) Immune response to <i>Neospora caninum</i> live tachyzoites in prepubertal female calves. Parasitol Res. 118 (10): 2945-55. Villa-Mancera, A. <i>et al.</i> (2021) Phage display-based vaccine with cathepsin L and excretory-secretory products mimotopes of <i>Fasciola hepatica</i>. induces protective cellular and humoral immune responses in sheep. Vet Parasitol. 289: 109340. Di Giacomo, S. <i>et al.</i> (2022) Assessment on Different Vaccine Formulation Parameters in the Protection against Heterologous Challenge with FMDV in Cattle. Viruses. 14 (8): 1781. Pooley, H.B. <i>et al.</i> (2019) The humoral immune response is essential for successful vaccine protection against paratuberculosis in sheep. BMC Vet Res. 15 (1): 223.
Storage	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>
Guarantee	12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10094 available at:
<https://www.bio-rad-antibodies.com/SDS/MCA2440P>
10094

Regulatory For research purposes only

Related Products

Recommended Useful Reagents

[AbGUARD® HRP STABILIZER PLUS \(BUF052A\)](#)

[AbGUARD® HRP STABILIZER PLUS \(BUF052B\)](#)

[AbGUARD® HRP STABILIZER PLUS \(BUF052C\)](#)

[TMB CORE \(BUF056A\)](#)

[TMB CORE+ \(BUF062A\)](#)

[TMB SIGNAL+ \(BUF054A\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M390601:210923'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)