

Datasheet: MCA2462

Description:	RAT ANTI MOUSE CD80
Specificity:	CD80
Other names:	B7-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	RM80
Isotype:	lgG2a
Quantity:	0.25 mg

#### **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				1/10 - 1/50
Immunohistology - Frozen				
Immunohistology - Paraffin				
ELISA				
Immunoprecipitation				
Western Blotting			•	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein C supernatant	G from tissue culture
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> )	

Carrier Free	Yes			
Approx. Protein Concentrations	IgG concentration 1.0mg/ml			
Immunogen	BCL1 cells expressing CD80.			
External Database Links	UniProt:  Q00609 Related reagents			
	Entrez Gene:  12519 Cd80 Related reagents			
Synonyms	B7			
RRID	AB_905981			
Fusion Partners	Spleen cells from immunised SD rats were fused with cells of the P3U1 myeloma cell line.			
Specificity	Rat anti Mouse CD80 antibody, clone RM80 recognizes mouse CD80 (B7-1), a ~60 kDa cell surface glycoprotein which is a member of the CD28/B7 family. In mice, CD80 is expressed on monocytes, peritoneal macrophages and dendritic cells, and expression may be significantly increased upon B lymphocytes by LPS and by IL-4.  CD80 has been identified as a ligand for CD28 and cytotoxic T-lymphocyte antigen-4 (CTLA-4), two structurally similar molecules expressed on T cells. CD28 and CTLA4 are two receptors that have essential but opposing functions in T-cell stimulation. The Interaction of CD80 with CD28 stimulates and sustains T cell responses, whereas the interaction of CD80 with CTLA4 is reported to inhibit T-cell responses.			
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.			
References	<ol> <li>Nuriya, S. <i>et al.</i> (1996) The differential role of CD86 and CD80 co-stimulatory molecules in the induction and the effector phases of contact hypersensitivity. Int Immunol. 8 (6): 917-26.</li> <li>Jin LP <i>et al.</i> (2004) Adoptive transfer of paternal antigen-hyporesponsive T cells induces maternal tolerance to the allogeneic fetus in abortion-prone matings. J Immunol. 173 (6): 3612-9.</li> <li>Nakajima, A. <i>et al.</i> (1997) Requirement of CD28-CD86 co-stimulation in the interaction between antigen-primed T helper type 2 and B cells. Int Immunol. 9 (5): 637-44.</li> <li>Nozawa, K. <i>et al.</i> (2001) Preferential blockade of CD8(+) T cell responses by administration of anti-CD137 ligand monoclonal antibody results in differential effect on development of murine acute and chronic graft-versus-host diseases. J Immunol. 167 (9): 4981-6.</li> <li>Bedoret, D. <i>et al.</i> (2009) Lung interstitial macrophages alter dendritic cell functions to prevent airway allergy in mice. J Clin Invest. 119 (12): 3723-38.</li> </ol>			

6. Legutko, A. et al. (2011) Sirtuin 1 Promotes Th2 Responses and Airway Allergy by

Repressing Peroxisome Proliferator-Activated Receptor-{gamma} Activity in Dendritic Cells. J Immunol. 187: 4517-29.

- 7. Deng, J. *et al.* (2010) Dendritic cells loaded with ultrasound-ablated tumour induce in vivo specific antitumour immune responses. <u>Ultrasound Med Biol. 36 (3): 441-8.</u>
- 8. Kim, I. *et al.* (2016) Immunological characterization of de novo and recall alloantibody suppression by CTLA4Ig in a mouse model of allosensitization. <u>Transpl Immunol. 38:</u> 84-92.
- 9. He, Y.T. *et al.* (2016) *In vitro* generation of cytotoxic T lymphocyte response using dendritic cell immunotherapy in osteosarcoma. Oncol Lett. 12 (2): 1101-6.
- 10. Curina, G. *et al.* (2018) Evaluation of immune responses in mice and sheep inoculated with a live attenuated *Brucella melitensis*. REV1 vaccine produced in bioreactor. <u>Vet Immunol Immunopathol</u>. 198: 44-53.
- 11. Inada, T. *et al.* (2009) Vaccines using dendritic cells, differentiated with propofol, enhance antitumor immunity in mice. Immunopharmacol Immunotoxicol. 31 (1): 150-7.

#### Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</a>
Regulatory	For research purposes only

#### Related Products

#### **Recommended Secondary Antibodies**

Goat Anti Rat IgG (STAR69...) FITC

Goat Anti Rat IgG (STAR73...) RPE

Rabbit Anti Rat IgG (STAR16...) <u>DyLight®800</u>

Rabbit Anti Rat IgG (STAR21...) HRP
Rabbit Anti Rat IgG (STAR17...) FITC

Goat Anti Rat IgG (STAR131...) Alk. Phos., Biotin

Goat Anti Rat IgG (STAR72...) HRP

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) DyLight®650, DyLight®800

# **Recommended Negative Controls**

Fax: +1 919 878 3751

RAT IgG2a NEGATIVE CONTROL (MCA1212)

North & South Tel: +1 800 265 7376

America

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_us@bio-rad.com

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

## Printed on 23 Jun 2022

© 2022 Bio-Rad Laboratories Inc | Legal | Imprint