

Datasheet: MCA2462A488

**BATCH NUMBER 158418**

<b>Description:</b>	RAT ANTI MOUSE CD80:Alexa Fluor® 488
<b>Specificity:</b>	CD80
<b>Other names:</b>	B7-1
<b>Format:</b>	ALEXA FLUOR® 488
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	RM80
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	100 TESTS/1ml

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/5

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse						
<b>Product Form</b>	Purified IgG conjugated to Alexa Fluor® 488 - liquid						
<b>Max Ex/Em</b>	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>Alexa Fluor®488</td> <td>495</td> <td>519</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	Alexa Fluor®488	495	519
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
Alexa Fluor®488	495	519					
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant						
<b>Buffer Solution</b>	Phosphate buffered saline						
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> ) 1% Bovine Serum Albumin						
<b>Approx. Protein</b>	IgG concentration 0.05mg/ml						

## Concentrations

---

**Immunogen** BCL1 cells expressing CD80.

---

## External Database Links

### UniProt:

[Q00609](#)   [Related reagents](#)

### Entrez Gene:

[12519](#) Cd80   [Related reagents](#)

---

**Synonyms** B7

---

**RRID** AB\_905981

---

**Fusion Partners** Spleen cells from immunized SD rats were fused with cells of the P3U1 myeloma cell line.

---

**Specificity** **Rat anti Mouse CD80 antibody, clone RM80** recognizes mouse CD80 (B7-1), a ~60 kDa cell surface glycoprotein which is a member of the CD28/B7 family. In mice, CD80 is expressed on monocytes, peritoneal macrophages and dendritic cells, and expression may be significantly increased upon B lymphocytes by LPS and by IL-4.

CD80 has been identified as a ligand for CD28 and cytotoxic T-lymphocyte antigen-4 (CTLA-4), two structurally similar molecules expressed on T cells. CD28 and CTLA4 are two receptors that have essential but opposing functions in T-cell stimulation. The interaction of CD80 with CD28 stimulates and sustains T cell responses, whereas the interaction of CD80 with CTLA4 is reported to inhibit T-cell responses.

---

**Flow Cytometry** Use 10ul of the suggested working dilution to label  $1 \times 10^6$  cells in 100ul.

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR ([BUF041A/B](#)).

---

## References

1. Nakajima, A. *et al.* (1997) Requirement of CD28-CD86 co-stimulation in the interaction between antigen-primed T helper type 2 and B cells. [Int Immunol. 9 \(5\): 637-44.](#)
2. Nozawa, K. *et al.* (2001) Preferential blockade of CD8(+) T cell responses by administration of anti-CD137 ligand monoclonal antibody results in differential effect on development of murine acute and chronic graft-versus-host diseases. [J Immunol. 167 \(9\): 4981-6.](#)
3. Jin LP *et al.* (2004) Adoptive transfer of paternal antigen-hyporesponsive T cells induces maternal tolerance to the allogeneic fetus in abortion-prone matings. [J Immunol. 173 \(6\): 3612-9.](#)
4. Inada, T. *et al.* (2009) Vaccines using dendritic cells, differentiated with propofol, enhance antitumor immunity in mice. [Immunopharmacol Immunotoxicol. 31 \(1\): 150-7.](#)
5. Bedoret, D. *et al.* (2009) Lung interstitial macrophages alter dendritic cell functions to prevent airway allergy in mice. [J Clin Invest. 119 \(12\): 3723-38.](#)
6. Deng, J. *et al.* (2010) Dendritic cells loaded with ultrasound-ablated tumour induce in

- vivo specific antitumour immune responses. [Ultrasound Med Biol. 36 \(3\): 441-8.](#)
7. Legutko, A. *et al.* (2011) Sirtuin 1 Promotes Th2 Responses and Airway Allergy by Repressing Peroxisome Proliferator-Activated Receptor- $\gamma$  Activity in Dendritic Cells. [J Immunol. 187: 4517-29.](#)
8. He, Y.T. *et al.* (2016) *In vitro* generation of cytotoxic T lymphocyte response using dendritic cell immunotherapy in osteosarcoma. [Oncol Lett. 12 \(2\): 1101-6.](#)
9. Kim, I. *et al.* (2016) Immunological characterization of de novo and recall alloantibody suppression by CTLA4Ig in a mouse model of allosensitization. [Transpl Immunol. 38: 84-92.](#)
10. Curina, G. *et al.* (2018) Evaluation of immune responses in mice and sheep inoculated with a live attenuated *Brucella melitensis*. REV1 vaccine produced in bioreactor. [Vet Immunol Immunopathol. 198: 44-53.](#)

---

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

---

**Guarantee** 12 months from date of despatch

---

**Acknowledgements** This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or [outlicensing@thermofisher.com](mailto:outlicensing@thermofisher.com)

---

**Health And Safety Information** Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2462A488>  
10041

---

**Regulatory** For research purposes only

## Related Products

### Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:Alexa Fluor® 488 \(MCA1212A488\)](#)

### Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

**North & South** Tel: +1 800 265 7376

**Worldwide** Tel: +44 (0)1865 852 700

**Europe** Tel: +49 (0) 89 8090 95 21

**America** Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M385415:210513'

**Printed on 29 Apr 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)