

Datasheet: MCA2462A647T

BATCH NUMBER 151603

Description:	RAT ANTI MOUSE CD80:Alexa Fluor® 647
Specificity:	CD80
Other names:	B7-1
Format:	ALEXA FLUOR® 647
Product Type:	Monoclonal Antibody
Clone:	RM80
Isotype:	IgG2a
Quantity:	25 TESTS/0.25ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/5

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse						
Product Form	Purified IgG conjugated to Alexa Fluor® 647 - liquid						
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>Alexa Fluor®647</td> <td>650</td> <td>665</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	Alexa Fluor®647	650	665
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
Alexa Fluor®647	650	665					
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin						
Approx. Protein	IgG concentration 0.05mg/ml						

Concentrations

Immunogen BCL1 cells expressing CD80.

External Database Links

UniProt:

[Q00609](#) [Related reagents](#)

Entrez Gene:

[12519](#) Cd80 [Related reagents](#)

Synonyms B7

RRID AB_2076010

Fusion Partners Spleen cells from immunised SD rats were fused with cells of the P3U1 myeloma cell line.

Specificity **Rat anti Mouse CD80 antibody, clone RM80** recognizes mouse CD80 (B7-1), a ~60 kDa cell surface glycoprotein which is a member of the CD28/B7 family. In mice, CD80 is expressed on monocytes, peritoneal macrophages and dendritic cells, and expression may be significantly increased upon B lymphocytes by LPS and by IL-4.

CD80 has been identified as a ligand for CD28 and cytotoxic T-lymphocyte antigen-4 (CTLA-4), two structurally similar molecules expressed on T cells. CD28 and CTLA4 are two receptors that have essential but opposing functions in T-cell stimulation. The interaction of CD80 with CD28 stimulates and sustains T cell responses, whereas the interaction of CD80 with CTLA4 is reported to inhibit T-cell responses.

Flow Cytometry Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR ([BUF041A/B](#)).

References

1. Nuriya, S. *et al.* (1996) The differential role of CD86 and CD80 co-stimulatory molecules in the induction and the effector phases of contact hypersensitivity. [Int Immunol. 8 \(6\): 917-26.](#)
2. Jin LP *et al.* (2004) Adoptive transfer of paternal antigen-hyporesponsive T cells induces maternal tolerance to the allogeneic fetus in abortion-prone matings. [J Immunol. 173 \(6\): 3612-9.](#)
3. Nakajima, A. *et al.* (1997) Requirement of CD28-CD86 co-stimulation in the interaction between antigen-primed T helper type 2 and B cells. [Int Immunol. 9 \(5\): 637-44.](#)
4. Nozawa, K. *et al.* (2001) Preferential blockade of CD8(+) T cell responses by administration of anti-CD137 ligand monoclonal antibody results in differential effect on development of murine acute and chronic graft-versus-host diseases. [J Immunol. 167 \(9\): 4981-6.](#)
5. Bedoret, D. *et al.* (2009) Lung interstitial macrophages alter dendritic cell functions to prevent airway allergy in mice. [J Clin Invest. 119 \(12\): 3723-38.](#)

6. Legutko, A. *et al.* (2011) Sirtuin 1 Promotes Th2 Responses and Airway Allergy by Repressing Peroxisome Proliferator-Activated Receptor- γ Activity in Dendritic Cells. [J Immunol. 187: 4517-29.](#)

Storage

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Acknowledgements

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information

Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2462A647T>
10041

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:Alexa Fluor® 647 \(MCA1212A647\)](#)

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M366968:200529'

Printed on 09 Feb 2024