

Datasheet: MCA2462SBUV400

Description:	RAT ANTI MOUSE CD80:StarBright UltraViolet 400
Specificity:	CD80
Other names:	B7-1
Format:	StarBright UltraViolet 400
Product Type:	Monoclonal Antibody
Clone:	RM80
Isotype:	IgG2a
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) Staining at lower temperatures generally requires longer incubation to reach antibody binding equilibrium, and this effect is more pronounced for StarBright Dyes due to their larger size. For optimum staining, Bio-Rad recommends incubating samples either overnight at 4°C or 1 hour at room temperature.

Target Species	Mouse		
Product Form	Purified IgG conjugated to StarBright UltraViolet 400 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	StarBright UltraViolet 400	335	394
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin 0.1% Pluronic F68 0.1% PEG 3350 0.05% Tween 20
Approx. Protein Concentrations	For information on the concentration of our StarBright Dye conjugated reagents please visit our FAQ page.
Immunogen	BCL1 cells expressing CD80.
External Database Links	UniProt: Q00609 Related reagents Entrez Gene: 12519 Cd80 Related reagents
Synonyms	B7
Fusion Partners	Spleen cells from immunized SD rats were fused with cells of the P3U1 myeloma cell line.
Specificity	Rat anti Mouse CD80 antibody, clone RM80 recognizes mouse CD80 (B7-1), a ~60 kDa cell surface glycoprotein which is a member of the CD28/B7 family. In mice, CD80 is expressed on monocytes, peritoneal macrophages and dendritic cells, and expression may be significantly increased upon B lymphocytes by LPS and by IL-4. CD80 has been identified as a ligand for CD28 and cytotoxic T-lymphocyte antigen-4 (CTLA-4), two structurally similar molecules expressed on T cells. CD28 and CTLA4 are two receptors that have essential but opposing functions in T-cell stimulation. The interaction of CD80 with CD28 stimulates and sustains T cell responses, whereas the interaction of CD80 with CTLA4 is reported to inhibit T-cell responses.
Flow Cytometry	Use 5µl of the suggested working dilution to label 0.5x10 ⁶ cells in 100µl. Best practices suggest a 5 min centrifugation at 6,000g prior to sample application.
References	1. Nakajima, A. <i>et al.</i> (1997) Requirement of CD28-CD86 co-stimulation in the interaction between antigen-primed T helper type 2 and B cells. Int Immunol. 9 (5): 637-44. 2. Nozawa, K. <i>et al.</i> (2001) Preferential blockade of CD8(+) T cell responses by administration of anti-CD137 ligand monoclonal antibody results in differential effect on development of murine acute and chronic graft-versus-host diseases. J Immunol. 167 (9): 4981-6. 3. Jin LP <i>et al.</i> (2004) Adoptive transfer of paternal antigen-hyporesponsive T cells induces maternal tolerance to the allogeneic fetus in abortion-prone matings. J Immunol. 173 (6): 3612-9. 4. Inada, T. <i>et al.</i> (2009) Vaccines using dendritic cells, differentiated with propofol, enhance antitumor immunity in mice. Immunopharmacol Immunotoxicol. 31 (1): 150-7. 5. Bedoret, D. <i>et al.</i> (2009) Lung interstitial macrophages alter dendritic cell functions to

- prevent airway allergy in mice. [J Clin Invest. 119 \(12\): 3723-38.](#)
6. Deng, J. *et al.* (2010) Dendritic cells loaded with ultrasound-ablated tumour induce in vivo specific antitumour immune responses. [Ultrasound Med Biol. 36 \(3\): 441-8.](#)
7. Legutko, A. *et al.* (2011) Sirtuin 1 Promotes Th2 Responses and Airway Allergy by Repressing Peroxisome Proliferator-Activated Receptor- γ Activity in Dendritic Cells. [J Immunol. 187: 4517-29.](#)
8. He, Y.T. *et al.* (2016) *In vitro* generation of cytotoxic T lymphocyte response using dendritic cell immunotherapy in osteosarcoma. [Oncol Lett. 12 \(2\): 1101-6.](#)
9. Kim, I. *et al.* (2016) Immunological characterization of de novo and recall alloantibody suppression by CTLA4Ig in a mouse model of allosensitization. [Transpl Immunol. 38: 84-92.](#)
10. Curina, G. *et al.* (2018) Evaluation of immune responses in mice and sheep inoculated with a live attenuated *Brucella melitensis*. REV1 vaccine produced in bioreactor. [Vet Immunol Immunopathol. 198: 44-53.](#)

Storage	This product is shipped at ambient temperature. Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA2462SBUV400
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M452207:260710'

Printed on 20 Aug 2026