

Datasheet: MCA2462SBV610

BATCH NUMBER 100008129

Description:	RAT ANTI MOUSE CD80:StarBright Violet 610		
Specificity:	CD80		
Other names:	B7-1		
Format:	StarBright Violet 610		
Product Type:	Monoclonal Antibody		
Clone:	RM80		
Isotype:	lgG2a		
Quantity:	100 TESTS/0.5ml		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugat	ed to StarBright Violet	610 - liquid
lax Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm
	StarBright Violet 610	403	607
eparation	Purified IgG prepared supernatant	I by affinity chromatog	raphy on Protein G
ffer Solution	Phosphate buffered s	aline	
eservative	0.09% Sodium Azide	(NaN ₃)	
abilisers	1% Bovine Serum All	oumin	
	0.1% Pluronic F68		
	0.1% PEG 3350		

Immunogen

BCL1 cells expressing CD80.

External Database Links

UniProt:

Q00609 Related reagents

Entrez Gene:

12519 Cd80 Related reagents

Synonyms

B7

Fusion Partners

Spleen cells from immunized SD rats were fused with cells of the P3U1 myeloma cell line.

Specificity

Rat anti Mouse CD80 antibody, clone RM80 recognizes mouse CD80 (B7-1), a ~60 kDa cell surface glycoprotein which is a member of the CD28/B7 family. In mice, CD80 is expressed on monocytes, peritoneal macrophages and dendritic cells, and expression may be significantly increased upon B lymphocytes by LPS and by IL-4.

CD80 has been identified as a ligand for CD28 and cytotoxic T-lymphocyte antigen-4 (CTLA-4), two structurally similar molecules expressed on T cells. CD28 and CTLA4 are two receptors that have essential but opposing functions in T-cell stimulation. The Interaction of CD80 with CD28 stimulates and sustains T cell responses, whereas the interaction of CD80 with CTLA4 is reported to inhibit T-cell responses.

Flow Cytometry

Use 5μ I of the suggested working dilution to label 10^6 cells in 100μ I. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

References

- 1. Nakajima, A. *et al.* (1997) Requirement of CD28-CD86 co-stimulation in the interaction between antigen-primed T helper type 2 and B cells. Int Immunol. 9 (5): 637-44.
- 2. Nozawa, K. *et al.* (2001) Preferential blockade of CD8(+) T cell responses by administration of anti-CD137 ligand monoclonal antibody results in differential effect on development of murine acute and chronic graft-versus-host diseases. <u>J Immunol. 167 (9):</u> 4981-6.
- 3. Jin LP *et al.* (2004) Adoptive transfer of paternal antigen-hyporesponsive T cells induces maternal tolerance to the allogeneic fetus in abortion-prone matings. <u>J Immunol.</u> 173 (6): 3612-9.
- 4. Inada, T. *et al.* (2009) Vaccines using dendritic cells, differentiated with propofol, enhance antitumor immunity in mice. Immunopharmacol Immunotoxicol. 31 (1): 150-7.
- 5. Bedoret, D. *et al.* (2009) Lung interstitial macrophages alter dendritic cell functions to prevent airway allergy in mice. <u>J Clin Invest.</u> 119 (12): 3723-38.
- 6. Deng, J. *et al.* (2010) Dendritic cells loaded with ultrasound-ablated tumour induce in vivo specific antitumour immune responses. Ultrasound Med Biol. 36 (3): 441-8.
- 7. Legutko, A. *et al.* (2011) Sirtuin 1 Promotes Th2 Responses and Airway Allergy by Repressing Peroxisome Proliferator-Activated Receptor-{gamma} Activity in Dendritic Cells. <u>J Immunol. 187: 4517-29.</u>
- 8. He, Y.T. et al. (2016) In vitro generation of cytotoxic T lymphocyte response using

dendritic cell immunotherapy in osteosarcoma. Oncol Lett. 12 (2): 1101-6.

9. Kim, I. et al. (2016) Immunological characterization of de novo and recall alloantibody suppression by CTLA4lg in a mouse model of allosensitization. Transpl Immunol. 38: 84-92.

10. Curina, G. et al. (2018) Evaluation of immune responses in mice and sheep inoculated with a live attenuated Brucella melitensis. REV1 vaccine produced in bioreactor. Vet Immunol Immunopathol. 198: 44-53.

Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA2462SBV610 20471
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

MOUSE SEROBLOCK FcR (BUF041A) MOUSE SEROBLOCK FcR (BUF041B)

North & South Tel: +1 800 265 7376 America

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M421794:230726'

Printed on 17 May 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint