

Datasheet: MCA2472B

Description:	MOUSE ANTI PHOSPHOTYROSINE:Biotin
Specificity:	PHOSPHOTYROSINE
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	PY20
Isotype:	IgG2b
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation			▪	
Western Blotting	▪			
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Chemical
Product Form	Purified IgG conjugated to Biotin - liquid
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.1% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml

Specificity

Mouse anti Phosphotyrosine antibody, clone PY20 recognizes phosphotyrosine, enabling the detection, characterisation and isolation of proteins containing phosphorylated tyrosine residues.

The phosphorylation of tyrosine acts as an important signal in the control of cell mitogenesis, differentiation, proliferation, and migration and occurs following the activation of intracellular tyrosine kinases, including the T-cell receptor (TCR), epidermal growth factor (EGF) and many families of receptor and non-receptor protein tyrosine kinases (PTKs), which catalyse the transfer of

ATP to a tyrosine residue on specific cell protein targets.

The binding of PY20 to phosphorylated tyrosines can be inhibited by free phosphotyrosine and phenylphosphate, but not by free phosphate, phosphoserine or phosphothreonine.

The affinity of PY20 for phosphotyrosine is 10^{-6} to 10^{-7} M.

References

1. Ruff-Jamison, S. *et al.* (1991) Heavy and light chain variable region sequences and antibody properties of anti-phosphotyrosine antibodies reveal both common and distinct features. [J Biol Chem. 266 \(10\): 6607-13.](#)
2. Takagi, S. *et al.* (1991) Intracellular localization of tyrosine kinase substrates beneath crosslinked surface immunoglobulins in B cells. [J Exp Med. 174 \(2\): 381-8.](#)
3. Glenney, J.R. Jr. *et al.* (1988) Monoclonal antibodies to phosphotyrosine. [J Immunol Methods. 109 \(2\): 277-85.](#)
4. Vendel, A.C. *et al.* (2009) B and T lymphocyte attenuator regulates B cell receptor signaling by targeting Syk and BLNK. [J Immunol. 182: 1509-17.](#)

Storage

Store at +4°C or at -20°C if preferred.
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M366997:200529'

Printed on 11 Aug 2020

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)