

Datasheet: MCA2476F

| Description:  | HAMSTER ANTI HUMAN EMR3:FITC |
|---------------|------------------------------|
| Specificity:  | EMR3                         |
| Format:       | FITC                         |
| Product Type: | Monoclonal Antibody          |
| Clone:        | 3D7                          |
| Isotype:      | lgG                          |
| Quantity:     | 0.1 mg                       |
|               |                              |

### **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

|                | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry |     |    |                | Neat - 1/10        |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

| Target Species                    | Human                                |                       |                         |               |
|-----------------------------------|--------------------------------------|-----------------------|-------------------------|---------------|
| Product Form                      | Purified IgG - liquid                |                       |                         |               |
| Max Ex/Em                         | Fluorophore                          | Excitation Max (nm)   | Emission Max (nm)       |               |
|                                   | FITC                                 | 490                   | 525                     |               |
| Preparation                       | Purified IgG prepared supernatant    | by affinity chromatog | raphy on Protein G from | tissue cultur |
| Buffer Solution                   | Phosphate buffered saline            |                       |                         |               |
| Preservative<br>Stabilisers       | 0.09% Sodium Azide                   |                       |                         |               |
| Approx. Protein<br>Concentrations | IgG concentration 0.1                | mg/ml                 |                         |               |
| Immunogen                         | ARHO-EMR3-CD97 (EGF1) transfectants. |                       |                         |               |

| External Database |  |
|-------------------|--|
| Links             | UniProt:   |
|                   | Q9BY15 Related reagents  |
|                   | Entrez Gene:   |
|                   | 84658 EMR3 Related reagents  |
| RRID              | AB_872002  |
| Fusion Partners   | Spleen cells from immunized Armenian hamster were fused with cells of the mouse SP2/0 myeloma cell line.   |
| Specificity       | Hamster anti Human EMR3 antibody, clone 3D7 recognizes human epidermal growth factor (EGF) module-containing mucin-like hormone receptor 3 (EMR3), a ~56 kDa member of the EGF-7 transmembrane (TM7) family of adhesion receptors. EMR3 is expressed at the cell surface as a heterodimer.   |
|                   | EMR3 is predominantly expressed on granulocytes, and at lower levels on mature myeloic cells, monocytes and dendritic cells. EMR3 is absent on lymphocytes, haematopoietic stem cells and myeloid progenitors. EMR3 is up-regulated during late stages of neutrophil differentiation and is a marker for terminally differentiated cells (Matmati et al.2007). |
| Flow Cytometry    | Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.  |
| References        | 1. Drewniak, A. <i>etal.</i> (2009) Changes in gene expression of granulocytes during <i>in vivo</i> granulocyte colony-stimulating factor/dexamethasone mobilization for transfusion purposes. <u>Blood. 113: 5979-98.</u>  |
|                   | 2. Gazendam, R.P. <i>et al.</i> (2016) Impaired killing of <i>Candida albicans.</i> by granulocytes mobilized for transfusion purposes: a role for granule components. <u>Haematologica. 101</u> (5): 587-96.  |
|                   | 3. van de Geer, A. <i>et al.</i> (2017) Characterization of buffy coat-derived granulocytes for clinical use: a comparison with granulocyte colony-stimulating factor/dexamethasone-   |
|                   | pretreated donor-derived products. Vox Sang. 112 (2): 173-82.  |
|                   | 4. Hansen, M. <i>et al.</i> (2018) Efficient production of erythroid, megakaryocytic and myeloid cells, using single cell-derived iPSC colony differentiation. <u>Stem Cell Res. 29: 232-44.</u>   |
| Further Reading   | 1. Carpenter, G. & Cohen, S. (1990) Epidermal growth factor. J Biol Chem. 265 (14): 7709-12.   |
| Storage           | This product is shipped at ambient temperature. It is recommended to aliquot and store at  |

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

#### Guarantee

12 months from date of despatch

**Health And Safety** Material Safety Datasheet documentation #10040 available at: Information https://www.bio-rad-antibodies.com/SDS/MCA2476F 10040 Regulatory For research purposes only

# **Related Products**

## **Recommended Negative Controls**

HAMSTER (ARMENIAN) IgG NEGATIVE CONTROL:FITC (MCA2356F)

## **Recommended Useful Reagents**

**HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)** 

North & South Tel: +1 800 265 7376 America

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Tel: +49 (0) 89 8090 95 21 Europe

Fax: +49 (0) 89 8090 95 50

Fax: +1 919 878 3751

Email: antibody\_sales\_us@bio-rad.com

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M385433:210513'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint