

Datasheet: MCA2484B

Description:	MOUSE ANTI HUMAN CD282:BIOTIN
Specificity:	CD282
Other names:	TLR2
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	11G7
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10
Immunohistology - Frozen			■	
Immunohistology - Paraffin			■	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG conjugated to Biotin - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1mg/ml
External Database Links	<p>UniProt:</p> <p>O60603 Related reagents</p> <p>Entrez Gene:</p> <p>7097 TLR2 Related reagents</p>

Synonyms	TIL4
Specificity	<p>Mouse anti Human CD282 antibody, clone 11G7 recognizes the human Toll-like receptor 2 (TLR2), a ~90 kDa type 1 transmembrane protein that is primarily expressed by peripheral blood monocytes. TLR2, also known as CD282, is reported to respond to a diverse range of bacterial cell wall components, mediating the innate immune response in co-operation with MD-2.</p> <p>Mouse anti Human CD282 antibody, clone 11G7 has been reported to block IL-6 cytokine production in response to ara-lipoarabinomannan (araLAM) and tripalmitoyl cysteinyl lipopeptide (Pam3CSK4) (Sandor et al. 2003). Clone 11G7 does not inhibit the production of cytokines induced by zymosan stimulation.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Sandor, F. <i>et al.</i> (2003) Importance of extra- and intracellular domains of TLR1 and TLR2 in NFkappa B signaling. J Cell Biol. 162 (6): 1099-110. 2. Zhou, S. <i>et al.</i> (2010) Discovery of a novel TLR2 signaling inhibitor with anti-viral activity. Antiviral Res. 87 (3): 295-306. 3. Royer, P.J. <i>et al.</i> (2010) The mannose receptor mediates the uptake of diverse native allergens by dendritic cells and determines allergen-induced T cell polarization through modulation of IDO activity. J Immunol. 185 (3): 1522-31. 4. Chen, E.S. <i>et al.</i> (2010) Serum amyloid A regulates granulomatous inflammation in sarcoidosis through Toll-like receptor-2. Am J Respir Crit Care Med. 181 (4): 360-73. 5. Nguyen, D.N. <i>et al.</i> (2016) Oral antibiotics increase blood neutrophil maturation and reduce bacteremia and necrotizing enterocolitis in the immediate postnatal period of preterm pigs. Innate Immun. 22 (1): 51-62. 6. Royer, P.J. <i>et al.</i> (2010) The mannose receptor mediates the uptake of diverse native allergens by dendritic cells and determines allergen-induced T cell polarization through modulation of IDO activity. J Immunol. 185 (3): 1522-31.
Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>Storage in frost-free freezers is not recommended.</p> <p>This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

Printed on 28 Jul 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)