

Datasheet: MCA2484PE

Description:	MOUSE ANTI HUMAN CD282:RPE
Specificity:	CD282
Other names:	TLR2
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	11G7
Isotype:	lgG1
Quantity:	100 TESTS

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .								
		Yes	No	Not Determined	Suggested Dilution				
	Flow Cytometry				Neat - 1/10				
	Where this product has not been tested for use in a particular technique this does not								
	necessarily exclude its a guide only. It is recor system using appropria	mmended that	at the use	r titrates the product f	g dilutions are given as or use in their own				
Target Species	Human								
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized								
Reconstitution	Reconstitute with 1ml distilled water								
Max Ex/Em	Fluorophore	Excitation N	lax (nm)	Emission Max (nm)					
	RPE 488nm laser	496		578					
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant								
Buffer Solution	Phosphate buffered sa	line							
Preservative Stabilisers	0.09% Sodium Azide(1% Bovine Serum A 5% Sucrose	-,							

External Database Links	UniProt: <u>O60603</u> <u>Related reagents</u> Entrez Gene: 7097 TLR2 Related reagents			
Synonyms	7097 TLR2 <u>Related reagents</u> TIL4			
RRID	AB_906132			
Specificity	Mouse anti Human CD282 antibody, clone 11G7 recognizes the human Toll-like receptor 2 (TLR2), a ~90 kDa type 1 transmembrane protein that is primarily expressed by peripheral blood monocytes. TLR2, also known as CD282, is reported to respond to a diverse range of bacterial cell wall components, mediating the innate immune response in co-operation with MD-2.			
	Mouse anti Human CD282 antibody, clone 11G7 has been reported to block IL-6 cytokine production in response to ara-lipoarabinomannan (araLAM) and tripalmitoyl cysteinyl lipopeptide (Pam3CSK4) (<u>Sandor <i>et al.</i> 2003</u>). Clone 11G7 does not inhibit the production of cytokines induced by zymosan stimulation.			
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.			
References	 Sandor, F. <i>et al.</i> (2003) Importance of extra- and intracellular domains of TLR1 and TLR2 in NFkappa B signaling. J Cell Biol. 162 (6): 1099-110. Royer, P.J. <i>et al.</i> (2010) The mannose receptor mediates the uptake of diverse native allergens by dendritic cells and determines allergen-induced T cell polarization through modulation of IDO activity. J Immunol. 185 (3): 1522-31. Chen, E.S. <i>et al.</i> (2010) Serum amyloid A regulates granulomatous inflammation in sarcoidosis through Toll-like receptor-2. Am J Respir Crit Care Med. 181 (4): 360-73. Royer, P.J. <i>et al.</i> (2010) The mannose receptor mediates the uptake of diverse native allergens by dendritic cells and determines allergen-induced T cell polarization through modulation of IDO activity. J Immunol. 185 (3): 1522-31. Royer, P.J. <i>et al.</i> (2010) The mannose receptor mediates the uptake of diverse native allergens by dendritic cells and determines allergen-induced T cell polarization through modulation of IDO activity. J Immunol. 185 (3): 1522-31. Zhou, S. <i>et al.</i> (2010) Discovery of a novel TLR2 signaling inhibitor with anti-viral activity. Antiviral Res. 87 (3): 295-306. Nguyen, D.N. <i>et al.</i> (2016) Oral antibiotics increase blood neutrophil maturation and reduce bacteremia and necrotizing enterocolitis in the immediate postnatal period of preterm pigs. Innate Immun. 22 (1): 51-62. 			
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.			
Guarantee	12 months from date of despatch			

Health A Informati	nd Safety on		afety Datasł <u>w.bio-rad-an</u>	e at:		
Regulato	ry	For resear	ch purposes			
Relate	d Product	S				
Recomm	nended Neg	gative Cor	ntrols			
MOUSE Ig	G1 NEGATIVI	E CONTROL	RPE (MCA9	<u>28PE)</u>		
Recom	nended Use	eful Reage	ents			
	EROBLOCK (I EROBLOCK (I					
North & South America	Tel: +1 800 265 7 Fax: +1 919 878		Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50
	Email: antibody_s	sales_us@bio-ra	id.com	Email: antibody_sales_uk@b	io-rad.com	Email: antibody_sales_de@bio-rad.com
To find a b	atch/lot specifi	c datasheet	for this produ	ict, please use our online	search tool at:	bio-rad-antibodies.com/datasheets

'M419469:230616'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint