# Datasheet: MCA2497 BATCH NUMBER 150608

Description:	MOUSE ANTI HUMAN HLA DP DQ DR
Specificity:	HLA DP DQ DR
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	Bu26
Isotype:	lgG1
Quantity:	0.2 mg

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further						
	information. For general	information. For general protocol recommendations, please visit www.bio-					
	rad-antibodies.com/protocols.						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry	•			1/100 - 1/200		
	Immunohistology - Frozen			•			
	Immunohistology - Paraffin			•			
	ELISA			•			
	Immunoprecipitation			•			
	Western Blotting			•			
	Where this product has n	ot been t	ested for	use in a particular tecl	nnique this does not		
	necessarily exclude its us	se in such	n procedu	res. Suggested workin	ng dilutions are given as		
	•	a guide only. It is recommended that the user titrates the product for use in their own					
	system using appropriate			•			
Target Species	Human						
Product Form	Purified IgG - liquid						
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> )						
Carrier Free	Yes						

Approx. Protein Concentrations	IgG concentration 1.0mg/ml		
RRID	AB_906085		
Specificity	<b>Mouse anti Human HLA DP DQ DR antibody, clone Bu26</b> reacts with DP, DQ and DR beta chains, which are expressed by antigen presenting cells, B cells, monocytes and activated T lymphocytes.		
	The major histocompatibility complex (MHC) is a cluster of genes that are important in the immune response to infections. In humans, this complex is referred to as the human leukocyte antigen (HLA) region. There are 3 major MHC class II proteins encoded by the HLA which are HLA DP, HLA DQ and HLA DR.		
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.		
References	<ol> <li>Chapple, M.R. <i>et al.</i> (1990) A phenotypic study of B lymphocyte subpopulations in human bone marrow. <u>Clin Exp Immunol. 81 (1): 166-72.</u></li> <li>Forte M <i>et al.</i> (1992) T-lymphocyte responses to <i>Pneumocystis carinii</i> in healthy and HIV-positive individuals. <u>J Acquir Immune Defic Syndr. 5 (4): 409-16.</u></li> <li>Knox, P.G. &amp; Young, L.S. (1995) Epstein-Barr virus infection of CR2-transfected epithelial cells reveals the presence of MHC class II on the virion. <u>Virology. 213 (1): 147-57.</u></li> <li>Hassell AB <i>et al.</i> (1992) MHC restriction of synovial fluid lymphocyte responses to the triggering organism in reactive arthritis. Absence of a class I-restricted response. <u>Clin Exp Immunol. 88 (3): 442-7.</u></li> <li>Kumararatne, D.S. <i>et al.</i> (1990) Specific lysis of mycobacterial antigen-bearing macrophages by class II MHC-restricted polyclonal T cell lines in healthy donors or patients with tuberculosis. <u>Clin Exp Immunol. 80 (3): 314-23.</u></li> <li>Shokri, F. <i>et al.</i> (1993) Immunophenotypic and idiotypic characterisation of the leukaemic B-cells from patients with prolymphocytic leukaemia: evidence for a selective expression of immunoglobulin variable region (IGV) gene products. <u>Leuk Res. 17 (8): 669-76.</u></li> <li>Pithie AD <i>et al.</i> (1992) Generation of cytolytic T cells in individuals infected by <i>Mycobacterium tuberculosis</i> and vaccinated with BCG. <u>Thorax. 47 (9): 695-701.</u></li> </ol>		
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.		
Guarantee	12 months from date of despatch		
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2497 10040		

## **Related Products**

### **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12)	RPE				
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>					
Goat Anti Mouse IgG (STAR76)	RPE				
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>				
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,				
	DyLight®650, DyLight®680, DyLight®800,				
	FITC, HRP				
Goat Anti Mouse IgG (STAR77)	HRP				
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>				
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP				
Rabbit Anti Mouse IgG (STAR13)	HRP				
<b>Recommended Negative Controls</b>					

### MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	ad.com	Email: antibody_sales_uk@bio-	rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M367052:200529'

### Printed on 14 Jun 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint