

Datasheet: MCA2502F

Description:	MOUSE ANTI HUMAN CD74:FITC
Specificity:	CD74
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	Bu45
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)				Neat - 1/5

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) Results maybe enhanced using membrane permeabilisation. Membrane permeabilization is required for this application. The use of Leucoperm (Product Code <u>BUF09</u>) is recommended for this purpose.

Target Species	Human			
Product Form	Purified IgG conjuga	ated to Fluorescein Isoth	niocyanate Isomer 1	(FITC) - li
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525	
Preparation	Purified IgG prepare supernatant	ed by affinity chromatog	raphy on Protein A	from tissue
Buffer Solution	Phosphate buffered	saline		
Buffer Solution Preservative	Phosphate buffered			

Approx. Protein Concentrations	IgG concentration 0.1mg/ml
Immunogen	B lymphoblastoid cell line HFB1.
External Database Links	UniProt: P04233 Related reagents Entrez Gene:
	972 CD74 Related reagents
Synonyms	DHLAG
RRID	AB_844513
Fusion Partners	Spleen cells from immunized mice were fused with cells of the Ag8.653 myeloma cell line.
Specificity	Mouse anti Human CD74 antibody, clone Bu45 recognizes human CD74, a type II transmembrane protein, also known as the MHC class II associated invariant chain. The Bu45 antibody recognizes 33, 35 and 41 kDa isoforms of CD74. The CD74 molecule plays a critical role in the presentation of peptides, by the MHC class II antigens, to CD4 positive lymphocytes.
	CD74 is expressed on MHC class II positive cells including B cells, a subset of activated T cells, monocytes, dendritic cells and by various types of carcinoma. Expression of CD74 is primarily intracellular with moderate expression at the cell surface of B cells and monocytes. Mouse anti Human CD74 antibody, clone Bu45 binds to an epitope within the extracytoplasmic domain of the CD74 molecule, which is distal from the C-terminus.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	1. Wraight, C.J. <i>et al.</i> (1990) Human major histocompatibility complex class II invariant chain is expressed on the cell surface. <u>J Biol Chem. 265 (10): 5787-92.</u> 2. van Bergen, J. <i>et al.</i> (1997) Efficient loading of HLA-DR with a T helper epitope by genetic exchange of CLIP. <u>Proc Natl Acad Sci U S A. 94: 7499-502.</u> 3. Beswick, E.J. <i>et al.</i> (2005) <i>Helicobacter pylori</i> binds to CD74 on gastric epithelial cells and stimulates interleukin-8 production. <u>Infect Immun. 73: 2736-43.</u> 4. Barrera, C.A. <i>et al.</i> (2005) Polarized expression of CD74 by gastric epithelial cells. <u>J Histochem Cytochem. 53: 1481-9.</u> 5. Beswick, E.J. <i>et al.</i> (2006) The <i>Helicobacter pylori</i> urease B subunit binds to CD74 on gastric epithelial cells and induces NF-kappaB activation and interleukin-8 production. <u>Infect Immun. 74: 1148-55.</u> 6. Lapaque, N. <i>et al.</i> (2009) The HLA-DRalpha chain is modified by polyubiquitination. <u>J Biol Chem. 284: 7007-16.</u> 7. Goldenberg, D.M. <i>et al.</i> (2019) Subcutaneous administration of anti-CD74 antibody for

systemic lupus erythematosus <u>US patent US 20170266281 A1</u>

Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA2502F 10041
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

Recommended Useful Reagents

LEUCOPERM (BUF09)
HUMAN SEROBLOCK (BUF070A)
HUMAN SEROBLOCK (BUF070B)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M404206:220820'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint