

# Datasheet: MCA2503PE

Description:	MOUSE ANTI HUMAN CD49d:RPE
Specificity:	CD49d
Other names:	INTEGRIN ALPHA 4 CHAIN, VLA-4
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	Bu49
lsotype:	lgG1
Quantity:	100 TESTS

## **Product Details**

Applications	This product has been reported to work in the following applications. This information derived from testing within our laboratories, peer-reviewed publications or personation communications from the originators. Please refer to references indicated for furth information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes No	Not Determined	Suggested Dilution		
	Flow Cytometry	•		Neat		
	Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.					
Target Species	Human					
Species Cross Reactivity	Reacts with: Dog <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.					
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized					
Reconstitution	Reconstitute with 1.0 ml distilled water Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.					
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)			
	RPE 488nm laser	496	578			

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> ) 1% Bovine Serum Albumin 5% Sucrose
Immunogen	Hyatt myeloma cell line
External Database Links	UniProt:         P13612       Related reagents         Entrez Gene:         3676       ITGA4         Related reagents
Synonyms	CD49D
Fusion Partners	Cells from immunised mice were fused with cells of the NS0 myeloma cell line
Specificity	<b>Mouse anti Human CD49d monoclonal antibody, clone Bu49</b> recognizes human CD49d, also known as integrin alpha 4 or VLA-4 alpha subunit. CD49d is a ~150 kDa single pass type I transmembrane glycoprotein. CD49d can be proteolytically cleaved to yield fragments of ~80 and ~70 kDa (Hemler <i>et al.</i> 1987). CD49d associates with either CD29 to form VLA-4 or with Integrin beta-7 to form The Peyer's patches-specific homing receptor LPAM-1, involved in the lymphocyte migration and homing to gut-associated lymphoid tissue (Sackstein 2006) through its interaction with MadCam-1, preferentially expressed on Peyer's patch high endothelial venules and postcapillary venules in lamina propria (Briskin <i>et al.</i> 1997).
	CD49d is expressed on a broad range of cells including lymphocytes, monocytes, eosinophils and thymocytes. CD49d expression may serve as a prognostic parker of survival and disease progression in patients with chronic lymphocytic leukemia and may have therapeutic potential ( <u>Bulian <i>et al.</i> 2008</u> ).
	Mouse anti human CD49, clone Bu49 cross-reacts with the canine B-cell leukemia cell line GL-1 and induces homotypic cell aggregation ( <u>lkewaki <i>et al.</i> 2010</u> ).
Flow Cytometry	Use 10ul of the suggested working dilution to label $1 \times 10^6$ cells in 100ul
References	<ol> <li>Oxford University Press. (1995) Leucocyte Typing V. White cell differentiation antigens. Volume Two. p 1617-8.</li> <li>Ikewaki, N. <i>et al.</i> (2010) Anti-human very late antigen-alpha4 (CD49d) monoclonal antibody (BU49) cross-reacts with the canine B-cell leukemia cell line GL-1, resulting in the induction of homotypic cell aggregation. <u>Cell Immunol. 263: 55-64.</u></li> <li>Hamilton, E. <i>et al.</i> (2012) Mimicking the tumour microenvironment: three different</li> </ol>

	co-culture systems induce a similar phenotype but distinct proliferative signals in primary chronic lymphocytic leukaemia cells. <u>Br J Haematol. 158: 589-99.</u>
Storage	Prior to reconstitution store at +4°C. After reconstitution store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA2503PE 20487
Regulatory	For research purposes only

## **Related Products**

### **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA928PE)

### **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	id.com	Email: antibody_sales_uk@bio-rad	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M419492:230616'

#### Printed on 19 Oct 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint